



# CARBON FARMING SCHEME

LIFE Preparatory Project



## **Draft report on calculation methods to be applied in estimating quantitatively agricultural and forest carbon sinks and their stability**



LIFE19 PRE FI001 – SI2.828588  
The Life Carbon Farming  
project has received funding  
from the LIFE Programme of  
the European Union

## LIFE Carbon Farming Scheme

# Expanding carbon sequestration activities by providing best practices and guidance for future farming schemes

The goal of the project is to identify and accelerate the development and adoption of novel incentives for carbon sequestration and the increase and maintenance of the organic carbon stock in soil and biomass in Europe. With the aim of promoting a well-functioning voluntary carbon market the project will uncover the key factors in supply and demand measures to invite the private sector to accelerate climate action. The results of the project will be fed into the development of the EU agricultural and climate policies.

[Read more: www.st1.com/st1-life](http://www.st1.com/st1-life)

### Coordinator:



### Partners:



Tyynelän  
tila



puro •  
earth

NEOT  
North European Oil Trade

### Financed by:



LIFE19 PRE FI001 – SI2.828588  
The Life Carbon Farming  
project has received funding  
from the LIFE Programme of  
the European Union

Draft report on calculation methods to be applied in estimating quantitatively agricultural and forest carbon sinks and their stability is part of the Action A1 work package lead by Natural Research Institute Finland.

Published 30.11.2020.

# LIFE CarbonFarmingScheme 30/11/2020

## Project Data

Project number: LIFE19 PRE FI/001 – SI2.828588

Project start date: 12/05/2020

Project end date: 11/05/2022

Total Project duration: 24 months

Project Website: <https://www.st1.com/st1-life>

## Beneficiary Data

Beneficiary: Natural Resources Institute Finland (Luke)

Contact person: Hannu Ilvesniemi

Postal address: Latokartanonkaari 9, FI-00790 HELSINKI

E-mail: [hannu.ilvesniemi@luke.fi](mailto:hannu.ilvesniemi@luke.fi)

# Contents

<b>1. Introduction</b>	<b>8</b>
<b>2. Research objectives</b>	<b>10</b>
<b>3. Background</b>	<b>10</b>
3.1. Carbon in terrestrial ecosystems	10
3.2. Carbon cycle of atmosphere-plant-soil ecosystems	12
3.3. Biomass carbon	14
3.4. Factors affecting biomass carbon pools	14
3.5. Soil organic carbon	15
3.6. Factors affecting soil organic carbon pools	16
3.7. Practices to promote carbon sequestration	18
<b>4. Detecting the change in the biomass and soil carbon pools</b>	<b>19</b>
4.1. Uncertainty quantification of measurements	22
<b>5. Carbon markets</b>	<b>23</b>
5.1. Issues associated with carbon sequestration projects	23
<b>6. Conventional methods</b>	<b>25</b>
6.1. Soil carbon stock determination	25
6.1.1 Sampling design and bulk density	26
6.1.2 Soil organic carbon content determination	27
6.1.3 Minimum number of sampling points	28
6.1.4 Sampling frequency	30
6.1.5 Uncertainties in soil sampling	30
6.1.6 Sampling cost	32
6.2. Biomass carbon stock estimation	32
6.2.1 National forest inventories	33
6.2.2 Allometric equations	33
6.2.3 Estimating carbon stock from forest inventories	35
6.2.4 Inventory planning and sample size	36
6.2.5 Uncertainties in forest inventories	37
6.2.6 Accuracy and cost	38
<b>7. Modelling soil carbon stocks</b>	<b>39</b>
7.1. Three levels of soil models	39

7.2. Soil carbon turnover models internal structure	43
7.3. SOC model scales	44
7.4. Model selection	45
7.5. Uncertainties and sensitivity analysis	46
7.6. Model performance	47
<b>8. Spectral methods</b>	<b>48</b>
8.1. Sensing soil organic carbon	49
8.2. Proximal soil sensing	50
8.2.1 Visible and infrared reflectance spectroscopy	51
8.2.2 Laser-induced breakdown and neutron induced gamma-ray spectroscopy	51
8.2.3 Uncertainties and important issues	52
8.2.4 Accuracy and cost	53
<b>9. Remote sensing</b>	<b>55</b>
9.1. Theory	55
9.2. Resolutions	56
9.3. Optical sensors	58
9.4. Synthetic aperture radar	59
9.5. Lidar	60
9.6. Important issues influencing biomass estimations for carbon mapping	61
9.6.1 Remotely sensed data	62
9.6.2 Reference data quality and variables	62
9.6.3 Modelling and uncertainty analysis	63
9.7. Accuracy	65
9.8. Cost	65
<b>10. Gas flux</b>	<b>66</b>
10.1. Eddy covariance	67
10.1.1 Important things to consider	68
10.1.2 Uncertainties and accuracy	69
<b>11. Other methodologies</b>	<b>70</b>
11.1. Verra	70
11.2. Gold Standard	72
<b>12. Calculation examples</b>	<b>73</b>
12.1. Forest stands	73
12.2. Agriculture	75

<b>13. Discussion</b>	<b>79</b>
13.1. Direct measurements	79
13.2. Spectral methods	80
13.3. Gas flux methods	80
13.4. Modeling methods	81
13.5. Concluding remarks	81
<b>14. References</b>	<b>82</b>

# PART I

# 1. Introduction

The global CO<sub>2</sub> balance in the atmosphere is depending on the balance between carbon inputs and outputs to atmosphere. Biomass accumulation is one of the available means to reduce carbon from atmosphere. In different EU countries there are several national tools to estimate the biomass production of plant species commonly used in agriculture and forestry. In this report we present a representative set of literature related to the processes relevant in estimating carbon stocks and stock changes in different types of ecosystems. The scope of the review was mainly in European situation, but also global issues have been taken into account. In the review we show the present understanding of the processes related to biomass carbon stocks and stock changes. The accuracy and representativity of the measured data and the effects of model structure and assumptions made in model calculations are also considered. This report will be the basis of more detailed analysis and suggestions for improvements to be presented in the final report of this project.

The literature review presented here is mainly based on the Ms. thesis of Anniina Lampinen to the University of Helsinki.

The natural carbon cycle is affected by human activity, and natural carbon pools have been decreasing. At the same time carbon concentration in the atmosphere and oceans has increased (Ciais et al. 2013). Disturbance of this natural cycle causes different negative impacts to current state of global climate and the states of the oceans. Carbon dioxide (CO<sub>2</sub>) is responsible for the majority of global warming, but methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O) and volatile organic compounds (VOCs) all are issues of concern (Toensmeier 2016). The concentration of these greenhouse gases (GHG) has increased in the atmosphere since industrialization. The use of fossil fuels as a source of energy, land use and changes in land use are the major causes of these rapidly elevated concentrations. CO<sub>2</sub> concentration has increased by 40% from 1750 to 2011, CH<sub>4</sub> 150% and N<sub>2</sub>O 20% in same time period (Ciais et al. 2013).

Land is needed for food production and living space and because population and per capita consumptions increases, demand for food and natural resources also grow continuously. This creates consequent stress to ecosystems. Global land use changes include for example deforestation and expansion of agriculture in tropics, afforestation and reforestation in temperate regions, intensification of agriculture and urbanization. The vegetation cover has been lost in many arid and semi-arid ecosystems in all climate domains (Song et al. 2018).

Terrestrial carbon locates naturally in soils and in above ground biomass. Carbon pool in soils is twice as large as that in atmosphere (Smith 2012). Thus, even small changes in this stock can influence the atmospheric CO<sub>2</sub> concentrations. Biomass carbon stock size is



approximately same as the atmospheric pool (Smith 2012). Carbon in soil and biomass can be released into the atmosphere due the burning of fossil carbon, land use changes, management practises or because of natural causes due the decomposing of organic matter (Janzen 2004). Soil carbon stocks are especially important because they can sequestrate large amounts of atmospheric CO<sub>2</sub>, which makes soil important factor to the global carbon balance (Bispo et al. 2017).

The Paris Agreement (2015) sets the target to limit climate change to 1.5°C. To reach that goal, all possible mitigation practises should be included into the global framework to avoid climate change. Carbon sequestration into natural pools could be a strategy for the removal of greenhouse gases from the atmosphere. Carbon sequestration is also called negative emission technology or carbon dioxide removal option (Smith 2016). Climate, land use, management and edaphic factors affect the amount of carbon stocks, but changes in those pools are not well understood. Without appropriate understanding it's hard to design monitoring, reporting and verification platforms (Smith et al. 2020). Carbon sequestration is also a reversible process so long-term monitoring is necessary to ensure that carbon sequestered persisted in these pools (Smith 2012). In this report only carbon sequestration through biological processes is considered.

Greenhouse gas emissions and carbon stocks are complex to measure. Climate, soil and vegetation characteristics and land management practises cause variation and large heterogeneity in carbon emissions and stocks. These factors vary largely in all spatial scales (Bispo et al. 2017). Terrestrial carbon pools have climate change mitigation potential with low associated costs (Elofsson and Gren 2018). To include these pools to EU climate policy, it would be necessary to quantify the carbon stock sizes and changes in stocks. For market confidence and to satisfy regulatory requirements the quantification methods should provide accurate results and at the same time being practical and financially achievable (Roxburgh et al. 2015). Measuring and monitoring of stocks is a key step towards sustainable carbon markets (Lankoski et al. 2020).

## 2. Research objectives


Purpose of this report is to gather information about currently used terrestrial carbon stock quantification methods. Under interest were especially uncertainties, accuracy, costs and scale associated to different methods. The aim was to compare the usability of different methods against developing carbon trading markets. Research questions are:

1. How can the size of biomass and soil carbon pools be determined?
2. Pros and cons of different carbon pool estimation methods as tool of carbon emission trading/carbon offsetting?
3. Example calculations of selected forest and agriculture farms

## 3. Background

### 3.1. Carbon in terrestrial ecosystems


There are five major carbon (C) pools in the Earth (Fig. 1): Ocean (77.4% of global carbon pool), fossil carbon (14.9%), soil (5%), biotic pool (1.2%) and atmospheric pool (1.5%). These pools are not constant but carbon cycles back and forth between them. Carbon cycle is a natural planetary cycle and it has occurred billions of years (Toensmeier 2016). Carbon's natural cycle has been affected by human actions: the amount of fossil carbon is decreasing annually due to the burning of fossil fuels, soil carbon pools has reduced since the start of industrialization because of land use management, and the same human activities have also decreased the size of the biotic pool. The atmospheric carbon increases because the other pools decrease, and the amount of carbon in oceans grow annually because it partially absorbs the excess carbon from the atmosphere (Toensmeier 2016).



**Fig. 1.** Illustrative picture of relative sizes of the different carbon pools on Earth.

Soil and biotic carbon are terrestrial carbon pools. Biotic carbon pool means living biomass, plants and animals, and dead detritus. Deforestation and agriculture have caused the loss of terrestrial carbon, the estimated loss amount is 320 billion tons of carbon, and majority of this has happened since 1850 (Toensmeier 2016). The excess carbon in the atmosphere causes global warming due to greenhouse gas effect, but soil degradation and loss of biotic pool have also other, complicated negative effects. Soil degradation have caused e.g. problems with soil fertility and erosion, especially in areas with intense soil use (Wild 1993). Loss of biodiversity is also well recognized problem (Dirzo and Raven 2003).

According to Jobbagy and Jackson (2000), the first three meters of mineral soils contain between 1500 and 2400 Pg of organic carbon. They estimated that first meter contains globally approximately 1500 Pg carbon, second and third meter 490 and 350 Pg carbon, respectively. Terrestrial vegetation contains approximately 450–650 Pg of carbon. Peat soils and permafrost account for more than 1500 Pg. These carbon pools are distributed across the terrestrial ecosystems (Fig. 2).




**Fig. 2.** Organic carbon pool sizes in different climatic regions. Green bars present above- and belowground biomass carbon. Topsoil (brown) and subsoil (orange) present the soil organic carbon pools. It's notable that terrestrial organic carbon pools are not distributed evenly across different climatic regions. Modified from Scharlemann et al. (2014), data is originally from Hiederer & Köchy (2011) and Ruesch and Gibbs by IPCC (2008).

## 3.2. Carbon cycle of atmosphere-plant-soil ecosystems

Terrestrial carbon pools interact constantly with atmospheric carbon via photosynthesis and respiration (Fig. 3). Through photosynthesis, plants convert sunlight, water and atmospheric carbon dioxide ( $\text{CO}_2$ ) into carbohydrates and oxygen. This photosynthesized carbon, which plants use for growth, creates the biomass carbon pool, or biomass carbon stock ( $\text{C}_{\text{biom}}$ ). According to Toensmeier (2016) Carbon that plants do not use directly can be transported to soil. Photosynthetic carbon is transported into the soil from plant roots as compounds that plant roots exude. There are more than 200 carbon-rich compounds that plants produce for different purposes. These exudates' roles include e.g. helping with nutrient cycling by feeding soil organisms, functioning as a suppressor of diseases, or to entice predators of pests (plant-microbe interaction in rhizosphere). Between 10 and 40% of photosynthesized

carbon passes through the roots within an hour (Toensmeier 2016). Over time, plants die, and aboveground biomass falls to the ground, where carbon-rich litter is partly decomposed. In this process, about 60–70% of carbon is released into the atmosphere as  $\text{CO}_2$ , and the remaining becomes soil organic matter (SOM) (Toensmeier 2016). Dead root biomass can become soil carbon as well. About half of soil organic matter is carbon (SOC). Respiration is the opposite reaction of photosynthesis and it describes the carbon flux from the soil to the atmosphere. Respiration can be divided into autotrophic respiration, which means carbon dioxide flux from plants, and heterotrophic respiration which refers to respiration of soil fauna. Rate of respiration and rate of photosynthesis depends from several environmental, climatic, soil characteristic and species-specific factors (Raich and Nadelhoffer 1989).

Carbon balance describes the balance of photosynthesis and plant and soil respiration (Fig. 3). If rate of photosynthesis is higher than total respiration rate, the ecosystem stores more carbon than it emits. These total fluxes determine, if the soil and biomass are carbon sinks and potential long-term carbon pools. Globally the annual flux of carbon between decomposition of organic matter and plant respiration is 119.7 Pg and photosynthesis flux is 123 Pg carbon per year, which makes soil a carbon sink (Bispo et al. 2017).



**Fig. 3.** Simplified chart picture presenting carbon cycle between atmosphere and terrestrial ecosystem. Arrows present the carbon transport between different carbon soil and biomass pools.

### 3.3. Biomass carbon

Plants sequester carbon within their biomass. Biomass carbon pools are aboveground and belowground biomass (IPCC 2006). Some of the synthesized glucose from photosynthesis is used for plant growth. Cellulose and lignin are important parts of plant cell wall and they are the backbone of plants providing structure and support. Cellulose and lignin have high carbon content and approximately 50% of the weight of the dehydrated plant biomass is carbon (Sedjo and Sohngen 2012).

### 3.4. Factors affecting biomass carbon pools

When plant sequesters carbon and produces glucose, it enables the growth, and larger growth means also higher photosynthesis rate. Net carbon uptake is highest in forests when the stands is closed and reasonably young (Sedjo and Sohngen 2012). Plant growth depends on temperature, radiation, moisture and nutrients and demand of these basic variables is species specific (Taiz and Zeiger 2010). In well balanced growth conditions, plant can achieve species specific maximum growth rate. Plant age, disturbances, climate and nutrient availability are the main controllers of plant productivity (Magnani et al. 2007). Nutrient availability is major the growth limiting factor in several ecosystems (LeBauer and Treseder 2008).

Because carbon sequestration is a reversible process, photosynthesized carbon can be released back to the atmosphere in respiration (Taiz and Zeiger 2010). Decomposition rate is highly dependent on temperature and moisture and it occurs mainly in soil (Lukac and Godbold 2011). Decomposition process is described with more detail in soil carbon section below.

Other important factors decreasing biomass carbon pools are harvesting, fires and herbivory (Sedjo and Sohngen 2012). Harvesting is the main factor affecting the amount of biomass carbon stocks. The manner of how harvested products are used determines the permanence of carbon. Harvested crops and trees that are used for short-lived products releases the carbon quickly back to the atmosphere. Trees can also be used to produce long-term products, which bind the carbon until wood starts to decay (Pukkala 2019). Fires (Fig. 4) and herbivory affect the amount of biomass. For example, in 2008, 189.7 Tg of CO<sub>2</sub> was released into the atmosphere from fires in US (Sedjo and Sohngen 2012).





**Fig. 4.** Boreal forest after a small wildfire. Ground was covered with few centimetres' depth layer of charred plant material and dead needles. Photo: Anniina Lampinen.

### 3.5. Soil organic carbon

Soil organic matter includes all organic components in the soil, and it is a complex mixture of compounds (Wild 1993). Litter is the first stage of organic matter entering the soil. Litter is dead plant material and it can originate from aboveground (canopy) or belowground (roots) (Lukac and Godbold 2011). Litter is full of nutrients and energy and their accessibility depends

on the quality of litter. Litter quality can be estimated with its ratio of total carbon to total nitrogen (C: N) (Lukac and Godbold 2011).

Soil organic matter carbon content is about 58% (Toensmeier 2016). Soil carbon can be divided into soil organic carbon (SOC) and soil inorganic carbon (SIC). Soil inorganic carbon comprehends mainly carbonate minerals calcite ( $\text{CaCO}_3$ ) and dolomite [ $\text{CaMg}(\text{CO}_3)_2$ ] and they are from geologic sources. Soil organic carbon is the carbon component of soil organic matter (SOM) (Lorenz and Lal 2018). According to Killham and Foster (1994) soil organic carbon can be separated into three pools or stocks: soluble, insoluble and biomass. Soluble fraction's decomposition rate is fast and because of that, soils consists only of 1% of soluble carbon. 90% of soil carbon is insoluble and it is a complex mixture of different plant materials in different decomposition states. Insoluble organic carbon forms the SOC stocks. Soil biomass (9%) comprehends soil microbes and animals which are responsible for most of the decomposition activity and carbon cycle (Killham and Foster 1994). Stable carbon is formed when carbon interact with soil particles (Killham and Foster 1994). Because soil organic matter comprehends a variety of different chemical compounds, it interacts with soil's mineral particles resulting in organo-mineral associations. Soil aggregates are one result from this interaction, and they are an important factor affecting soil carbon stability. Small humified compounds have high affinity for clay particles, and more than half of the total soil carbon is strongly bound to clay. This strong bond increases significantly the residence time of carbon in soils (Lukac and Godbold 2011).

### 3.6. Factors affecting soil organic carbon pools

Soil organic matter accumulation and distribution is affected by several biotic and abiotic factors and processes. Biotic factors include plant input and soil organisms. Important abiotic factors are e.g. climate (temperature and precipitation) and soil mineralogy (soil physio-chemical properties) (Luo et al. 2017). Anthropogenic factors have also an impact to soil carbon accumulation and distribution. Typical land management practises like fertilization with nitrogen and tillage affect to soil microorganism and structure (Jackson et al. 2017, Lorenz & Lal 2018). Soil organic carbon stocks are a result of complex interactions among several variables (Lukac and Godbold 2011).

Litter is mainly decomposed via biological processes. Soil microbes and other living organism produce enzymes and metabolic substances which drive the decomposition processes. Activity of those enzymes are temperature and moisture limited, and each have specific optimal range (Boyero et al. 2011).



Aboveground biomass sequesters carbon from the atmosphere and higher plant productivity often increases SOM in the soils. The relationship between net primary production (NPP) and SOC accumulation is not linear and bigger biomass does not automatically lead to increase in SOC pools (Jackson et al. 2017). Reasons for these nonlinear relationships are not well known because soil complex interactions and processes are not yet well understood. Some of the suggested reasons are soil carbon saturation (Mayzelle et al. 2014), the priming effect (Kuzyakov 2010) and carbon allocation between plant parts (Jobbagy and Jackson 2000). Soils have limited capacity to sequester carbon and they can saturate. The level of soil carbon saturation varies, and it's affected e.g. by soil mineralogy and climate (Mayzelle et al. 2014). The priming refers to a situation where plant input to soil increases microbial activity which leads to losses of accumulated SOM (Kuzyakov 2010). Carbon allocation between above- and belowground biomass can also be one of the factors explaining the nonlinear relationship between carbon input and SOC accumulation. Different plant species in diverse environments distribute carbon compounds differently between plant parts. The allocation of net primary production patterns varies between species and ecosystems, such ranging from 10% of carbon is allocated to roots in croplands, to 60% in native grasslands and 20% in forests (Poorter et al. 2012).

Carbon is not evenly spread in soil profile. Different soil layers typically include different amounts of carbon and this vertical variability is due the different decomposition rates of organic matter and their transportation (Jobbagy and Jackson 2000).

At the landscape scale, soil texture, pH, mineralogy, topology and land-use are the main factors affecting SOC heterogeneity. At the plot scale, plant species diversity and composition, and land management practises increase SOC heterogeneity (FAO 2019). Soil pH, clay content and cation exchange capacity all affect soil biochemical composition and distribution. These factors vary greatly on the ecosystem level but also on a smaller scale. Agricultural lands are relatively homogeneous compared to forests. But even in arable soils, where nitrogen deposit, pH and clay content are usually well monitored, spatial variation occurs (between parcels) on the farm level (Bispo et al. 2017).

Soil microbes affect stabilisation of carbon in soil. Soil microbes use carbon for growth and release it via respiration. When microbes die, these microbial residues or so called necromass, can be recycled as new substrates or it can be stabilized. Stabilization happens after necromass is bound to soil mineral surfaces and stored as microaggregates (Miltner et al. 2011). According to Kallenbach et al. (2016) and Balsler and Lian (2011) 50-80% of stable organic carbon in soils is necromass. Environmental and microbial controls, which are important factors controlling necromass recycling and thus soil carbon stabilization (Buckeridge et al. 2020).

## 3.7. Practices to promote carbon sequestration

Increasing the carbon input and decreasing the decomposition is the basic of carbon accumulation. Net primary production and decomposition rate varies between sites naturally, but there are management practises that are known to affect positively to the size of the different carbon stocks in forests and cultivated soils. Different agricultural and forestry practises e.g. residual management, soil tillage and fertilizer application add variation to carbon accumulation (Bispo et al. 2017). Global forest annual carbon sequestration potential is estimated to be 2-4 Gt C of atmospheric carbon. The “4 per mille Soils for Food Security and Climate” is an act which was launched to increase soil organic carbon content by 0.4% per year and at the same time to mitigate climate change. The 4p 1000 Initiative was a result from global survey where soil organic pool sizes and sequestration potentials were estimated. They reported that under best management practises 0.4% sequestration rate would be accomplished in cultivated soils in areas where topsoil carbon content is low, less than 30 t C/ha. With this rate, global agriculturals would be able to sequester 2-3 Gt carbon annually, which would offset 20-33% of anthropogenic emissions (Minasny et al. 2017).

In arable land, management practices and history of those affect the accumulation. Conservation practices have a technical potential to increase the soil carbon stocks (Lorenz & Lal 2018), and nowadays, term regenerative farming is commonly used term to describe those management practices that aim to minimize erosion and leaching of nutrients and organic matter, enhance biodiversity and soil health and to sequester carbon (Elevitch et al. 2018). Globally applied conservation principles include minimizing soil disturbance, maximizing surface cover and stimulate biological activity through cover crops, crop rotation and integrated nutrient and pest management (Lorenz & Lal 2018).

Tillage management, crop rotation, agroforestry, cover crops and application of organic amendments are land management practices that could increase carbon concentration in soils (Paustian et al. 2016). For example, tillage of soil produces more CO<sub>2</sub> efflux than soils that are under no-tilled soil management, because tillage makes soil organic material available to oxidation and microbial mineralization (Brekke et al. 2019). Selection of crop variate and avoiding the use of bare fallows have also positive impact to soil carbon accumulation (Lorenz & Lal 2018). Cultivation of cover crops increases the soil carbon accumulation by 32 g C/m<sup>2</sup> annually (Poeplau and Don 2015) and they also prevent nutrient leaching.

Afforestation is one way to increase the amount of biomass carbon. Tree and other vegetation growth in previously unforested sites would also increase the amount of carbon in soils (Sedjo and Sohngen 2012). Peltoniemi et al. (2004) studied how increasing stand age


would affect the soil carbon and they found out that carbon stock increased average  $4.7 \pm 1.4$  g/m<sup>2</sup>/year with increasing stand age. This accumulation was only noticed in organic layers, and no significant changes were measured in mineral soils.

Nitrogen fertilization in ecosystems that suffer from nitrogen deficiency is one possible way to affect the growth. For example, in boreal forests nitrogen fertilization increased growth significantly and in fertilization treated plots stemwood production increased by 29-37% compared to non-treated plots (Mäkipää et al. 1998). Nitrogen affects the growth, which means accumulation of carbon in biomass, but it also has impact to soil carbon pools. Whether the impact is negative, positive or neutral, depends on several factors. For example in boreal forests, soil organic matter accumulation is noticed to increase by nitrogen input (because of increase litter input) (Mäkipää 1995). Other forest management practises are thinning, extending the harvest rotation and selection of species varieties (breeding) (Sedjo and Sohngen 2012).

## 4. Detecting the change in the biomass and soil carbon pools

Carbon accumulation or losses can be in general determined in two different ways: measuring pool changes and measuring incoming and outgoing fluxes (Houghton 2003).

Estimations about different biomass and soil carbon pools are usually conducted with different methods (Fig. 5). Soil carbon pools can be quantified with direct soil sampling, sensing with spectroscopic methods or by modelling (Paustian et al. 2019). Biomass carbon pools can be quantified with inventory-based field measurements, remote sensing or modelling. Incoming and outgoing fluxes can be used to measure the whole ecosystem (biomass and soil) carbon pool size (Smith et al. 2020).




**Fig. 5.** Schematic picture of different carbon pool estimation methods and how they are linked together. Conventional soil and forest field samples provide information about current stock sizes and/or changes in those stocks. They also provided foundations to other methods (so called ground truthing). Measured values can be used to different kind of model development, initialization, validation etc. Spectral methods rely also to measured values, because there must be reference data to evaluate the reflectance. Modelling can be used to simulate carbon stock sizes or changes in those stocks. They also link the remotely sensed data to the ground truthing. Information from flux measurements and from remotely sensed data can be used to model development. Gas flux measurements present the net ecosystem carbon exchange.

Several European countries have launched networks to monitor the changes in soil condition (Morvan et al. 2008) and changes in biomass over time (Tomppo et al. 2010). But in European scale, the geographical distribution of monitoring networks is uneven, and in central and northern in Europe the number of sampling plots is higher than eastern Europe. Globally this same trend is recognized, and certain areas have representative study networks and some lack those completely. One large problem is also that networks differ considerably in their sampling protocols, designs, plot locations and sampling frequency. Sampling networks vary also within the country because some have different networks for different land-use types, like arable and forest lands (Heikkinen 2016, Tomppo et al. 2010). Long-term experiments are needed for data concerning processes that affect soil and biomass carbon and how different management practices affect the pools. These long-term field measurements also provide the basis for the calibration and validation of different models (Körschens 2005, Saarsalmi et al. 2014 Heikkinen 2016).

According to Pearson et al. (2007), to produce credible and transparent estimates of changes in include description of boundaries, project area, number of sample plots, project duration and monitoring frequency, 2. Information about the number of samples and other sampling protocols, 3. Carbon stock estimation methods and analyzing of the results. What

are the actual methods and how they are utilized? What techniques are used for result analyzing? 4. Net change estimation. How the change in carbon stocks can be estimated? 5. A quality control plan. What is the accuracy in these estimations? Quality assurance (QA) and quality control (QC) for results (Fig. 6).carbon pools the following steps are needed: 1. A monitoring plan should



**Fig. 6.** Steps needed for estimation of changes in carbon pools.

According to Pearson et al. (2007), carbon stock estimations should be primary conducted with field measurements, because of their accuracy. Pearson et al. (2007) also suggested that precision target should be that with 95% confidence level, the true population value is  $\pm 10\%$  of the sample estimate. Because field measurements are not the most cost-effective way to monitor carbon stock changes, alternative methods could be used, if those reach robust precision enough. Good common practices for carbon accounting secure the reliable results. Good practices according to Watson (2009) are summarize in table (Table 2).

**Table 2.** Important factors that should be considered in carbon pool estimations.

<b>Accurate and precise</b>	Both accuracy and precision should be achieved. Biases and uncertainties should be removed as far as it possible.
<b>Comparable</b>	Assumptions, methods and data must be commonly accepted (scientific consensus) and should provide meaningful and valid results between areas.
<b>Complete</b>	All relevant carbon pools should be included. If some are excluded, it should be well justified and documented.
<b>Consistent</b>	Estimates from different quantifications should present the actual difference between pools. Differences should not emerge from differences in methods.
<b>Relevance</b>	Trade-offs between time, resources, data and methods should be appropriate to the purpose of the quantification.
<b>Transparent</b>	Results should be able to be confirmed by a third party.

## 4.1. Uncertainty quantification of measurements

All carbon stock and stock change estimation methods include sources of uncertainties or errors and there is no such method which would provide absolute true pool size value. It's important to identify the sources of errors and quantify their nature and magnitude. Estimations of these uncertainties are especially important if the results are used for policy purposes. For example, sustainable creditable carbon unit is a result of quantification method which full uncertainty is defined and considered. Resolution is also an important term. Resolution can be defined as the smallest change in the measured quantity that instrument can detect (Aubinet et al. 2012).

Precision, accuracy, and bias are commonly used terms when discussing about uncertainties related to measurements. Precision refers to the degree of agreement in a series of measurements and accuracy is the closeness of a measurement to the true value (Husch et al. 2003). Bias is a term that describes the difference between the true value of a target and its average measured value. Bias error can also be called systematic error. Systematic errors are consistent and repeatable errors. Random errors can be called for precision errors and those types lack repeatability. Scattered result in a repeated measurement is a common sign of random error (Husch et al. 2003). These two traditional error types propagate in different ways and thus it is important to differ those. Random errors are impossible to correct due to their random nature, and they typically cause noise and scatter in the data. Random errors reduce the precision of the measurements. Repeated measurements (increasing the  $n$ ) is the only way to characterize the total random error. Averaging over  $n$  measurements improves the precision and result of this gives the standard error of the mean. Systematic error stays constant and it cannot be identified through statistical analysis (bias) (Aubinet et al. 2012).

Many carbon estimation methods rely on data-model fusion and different error types affect differently to the models (Lasslop et al. 2008). All models are as good as their most inaccurate parameter. Different error types at different stages affect the total uncertainty of the estimation. It's important to evaluate the magnitude of all possible errors.

# 5. Carbon markets

The Paris Agreement was launched 2015 and parties of the UNFCCC agreed to fight against climate change and to enhance investments needed for reach low carbon future. The Agreements central aim is to limit the temperature rise to 1.5 °C and thus prevent the harmful consequences of climate change. The Paris Agreement considers carbon sinks and reservoirs also as important factor as a tool for mitigating climate change, and The Agreement encourages Parties to enhance pools and to increase their sizes (United Nations 2020). European aims to be first carbon neutral continent in 2050 (European Comission 2020). To meet this target the EU is set several action plans, and for example European Green Deal is a package which includes first climate action initiatives, like European Climate Law (European Comission 2020). European Climate Law is going to be a first law where EU aims to write the year 2050 climate neutral target. European Climate Law would ensure that all EU policies would make effort to reach that target. To reach that target all possible mitigation practices should be adopted and to enhance practices to increase natural-based carbon sequestration, political incentive should be launched (European Comission 2020). One possible way is international carbon markets and the Paris Agreement recognize the importance of this in Articla 6, where it's mentioned that carbon trading would help to achieve emission reduction targets. This kind of market mechanism would include that farmers and foresters could manage their land so that carbon sequestration is maximize and would achieve economic value from these practices by selling the creditable carbon units they have produce in the carbon pool. Creditable carbon unit is not yet defined, and there is no such regulated mechanism where transaction would be conducted.

## 5.1. Issues associated with carbon sequestration projects

Climate change mitigation by changing the land management practises and at the same time providing excess income possibilities for farmers and foresters is a great goal to achieve. This approach would be a way to utilize the already existing, and possibly quite effective methods, but there are also several well-known issues associated to carbon sequestration projects via natural methods. Project issues usually contain the criteria which should be met so that it is possible to say that carbon is sequestered from the atmosphere and it's now in adequately stable form and locates in stock. Baseline determination, additionality, possible carbon leakage and permanence (and non-permanence risk) are the most discussed issues (García-Oliva 2004, Sedjo and Sohngen 2012).



In carbon market framework, the baseline determination is the first thing to address. Credible and accurate baseline is needed for monitoring difference in carbon stocks. Baseline determination is also one of the most challenging issues. According to Carcia-Olivia (2004) two approaches have been used for developing and applying baselines. One approach is project specific. Baseline is established case-by-case. Other is generic, where baseline is determined by using regional or national data. Important component of assessing carbon sequestration is also to determine whether the carbon benefits of an activity are truly additional. Carbon benefits of certain project/action should be compared to carbon stocks between a with- and without- action scenario. Leakage can occur when measurable carbon net change (decrease or increase) occurs outside wanted area due to project activity. For example, if farmer afforests land and after the afforestation of area deforests another land for agricultural purposes, the resulting carbon emissions are referred to as leakage. Monitoring and accounting leakage can be done either project specific or standardized. And because carbon sequestration is a reversibly process and carbon stored in terrestrial ecosystems are vulnerable to natural or anthropogenic disturbances. Fires and pests, harvesting and changes in land management and land-use may result to carbon release to the atmosphere (Carcia-Olivia 2004, Sedjo and Sohngen 2012).

Besides the tight criteria, there are also other challenges which should be resolved before well-functioning CO<sub>2</sub> emission markets. One big issue is high measuring, reporting and verifying (MRV) costs of carbon sequestration. Current costs are high and MRV processes are complicated and they lack standards. Small-scale projects are important for the development of local carbon markets and MRV costs should be reduced in order to allow the progress. High MRV costs are identified as a challenge and some solutions are also presented. For example, group certification options, more diverse group of auditors to carry out verification and baseline and stock change estimation with less time-consuming methods would reduce the costs. One approach is to value other benefits besides carbon to allow higher carbon prices (Köhl et al. 2020). Well implemented carbon sequestration projects are usually multi beneficial, and environmental and social benefits should be added to carbon price value. Higher carbon value would increase the project profitability. This is a possible approach, but it also adds uncertainties in the achievement evaluation state. It's hard and expensive to determine the size of the carbon stock change accurately and it would be even harder to evaluate the project impact for biodiversity.



## 6. Conventional methods

Sequestered carbon locates in aboveground and belowground stocks. The rate of carbon accumulation and the rate of carbon loss during a certain time period determines the stock size. The amount of carbon that can be stored in the biomass or soil is also an important defining factor affecting ultimate stock size (saturation) (Mayzelle et al. 2014, FAO 2019). Soil carbon stocks can be measured directly in units of carbon, but biomass is converted to units of carbon by multiplying biomass by 0.5 (IPCC default) or more specific values, if there is available data (Pearson et al. 2007).

Soil organic carbon stock estimation is challenging, because soil carbon content and soil biochemical processes vary greatly spatially and temporally. Physical sampling can be used for baseline determination and monitoring SOC stock changes. Soil sampling should be well planned and fulfil standard methodology criteria. Several environmental factors cause heterogeneity in the organic carbon content in soils and standardization adds confidence in sampling results (FAO 2019). For example 60% of carbon in boreal forests is located in soil (Liski et al. 2006) and changes in this pool are small in relation to the size of the stock, which means that forest soil carbon stock changes are hard to measure and detect reliably (Peltoniemi et al. 2004).

Aboveground vegetation can be measured with field work combined empirical models. Roots and other belowground biomass are more difficult to estimate, and usually belowground biomass is added to stock estimates with models without any actual sampling (Pearson et al. 2007).

### 6.1. Soil carbon stock determination

The conventional approach to determine soil organic carbon content is to collect soil samples and analyse them for carbon concentration. This procedure includes field soil sampling, sample preparation and laboratory analysis. Bulk density measurements are also required to convert concentration from mass to volume based (Whitehead et al. 2012). Soil dry bulk density is the dry mass per unit volume of the soil. Soils include solids and pores, and those pores can contain air, water or both and bulk density typically has high spatial variability (Wild 1993). Soil carbon concentration can be determined with dry combustion. Dry combustion is done with standard automatic instruments in a laboratory. Bulk density is conventionally measured with the volumetric ring method or the clod method (for soils with many rock fragments) (England and Viscarra Rossel 2018). Due to the soil characteristics

the direct soil sampling needs to be considered thoroughly in order to achieve effective and representative sampling (FAO 2019).

Soil sampling can be carried out in arable and forest soils, and every soil type. Forest soil sampling is more complicated than sampling of cultivated soils, but there are approaches that are suitable for soils under different vegetation types (grasslands, trees, wet peat etc.). Forest soils typically have well-developed organic layers and high organic matter content, which causes significant variations of SOC with depth. Compared to agricultural soils the spatial variability is usually larger in forest soils (IPCC 2000, Vanguelova et al. 2016).

### **6.1.1 Sampling design and bulk density**

Sampling size and core depth depend on the purpose of the sampling, and vertical and horizontal stratification or heterogeneity of soil carbon. If the purpose is to estimate the total amount of soil organic carbon stock in a certain area, then the sample collection should contain all subunits in that location (Whitehead et al. 2012). If the land area is shallow mineral soil with organic horizon, then the sampling of multiple cores should be distributed accordingly. Slopes also need several sampling points in different slope positions. In general that the larger the horizontal or vertical gradient, the larger the replicate needed. The horizontal gradient affects the number of cores and the vertical gradient affects the number of the samples collected in each core (Whitehead et al. 2012). Sampling locations for soil coring should be randomly determined to avoid bias. GPS coordinates of each sampling location should be recorded for future revisiting (FAO 2019).

Soil samples are collected from certain soil depth and most often soil sampling for carbon stock estimation purposes is done in a depth of 30 cm, the minimum recommended (FAO 2019). Deeper layers of soil can also be sampled in the depths of 30-60 cm and 60-100 cm. The 100 cm sampling depth often requires specific machinery (Smith et al. 2020). A sample can be collected by using a soil corer tool of known volume, or a pit. A large pit is more time consuming but reveals the whole soil profile and reduces uncertainties related to soil compaction. Soil coring with suitable tool and the pit are both accepted practices; most important is to use such methods that the needed parameters can be calculated/estimated (e.g. soil bulk density, soil mass) (FAO 2019).

Soil coring is usually conducted with a simple cylinder, which is pressed into the soil, to the depth of interest. FAO recommends that the core diameter is something between 50 and 100 mm. This is because a diameter smaller than 50 mm gives a small sample volume that makes it difficult to represent properly the coarse roots and coarse mineral fragments. Diameters larger than 100 mm are difficult to handle (FAO 2019). Depending on the depth of

the core, the sample might be divided into sections. According to the method from Blackmore et al. (1987), the fresh sample is weighted (each section if necessary), sub-sampled and homogenized. Dry mass is calculated from the water content and field weight, and bulk density is determined by dividing the dry mass by its volume (each section).

Compositing is a procedure where several soil cores (subsamples) are pooled together into one homogenous composite sample. This method is also known as bulking. In those samples, SOC concentration should be equal to the average SOC value of individual cores. Compositing method can be used for reducing spatial variability and overall cost from multiple soil sample analyses (FAO 2019).

## 6.1.2 Soil organic carbon content determination

Dry combustion is an analytical method to measure organic carbon content in soil. In dry combustion, finely grounded soil samples are burned generally around 1000 °C (Nelson and Sommers 1996). Pure oxygen acts as a catalyst or accelerator and ensures complete combustion of the sample. Other catalysts are vanadium pentoxide, copper (Cu), copper oxide (CuO) and aluminium oxide (Al<sub>2</sub>O<sub>3</sub>). The end product, CO<sub>2</sub>, is then quantified by gas chromatography. Since all carbon units are measured, it is important to remove other carbonates (SIC) before SOC determination. This is conducted with hydrochloric acid acidification prior the analysis. Some soils include high amounts of highly stable organic carbon compounds (e.g. char from natural fires and biochar) which don't decompose in temperatures under 600 °C. This may lead to underestimation of SOC concentration (Nelson and Sommers 1996). Other analytical methods are also available (e.g. wet digestion/oxidation) (Vitti et al. 2016.).

SOC content in a collected sample is then defined by measuring certain parameters and using those parameters in calculations of SOC stock. Soil fraction is sieved, and soil that passes 2 mm sieve and the organic carbon in it, is the internationally accepted definition of operational SOC (fine earth fraction). For SOC stock determination, soils fine earth and coarse mineral fraction, organic carbon concentration in the fine earth fraction and soil bulk density or fine earth mass are necessary parameters. Those parameters can be used in equation 1 to calculate SOC stock (FAO 2019).

$$SOC_i \text{ stock } \left( \frac{Mg C}{ha} \right) = OC_i \times BD_{fine i} \times (1 - vGi) \times t_i \times 0.1 \quad (1)$$

where

SOC<sub>i</sub> = soil organic carbon stock (in Mg C /ha) of the depth increment i

OC<sub>i</sub> = organic carbon content (mg C g/soil) of the soil fraction (< 2 mm) in the depth increment i

BD<sub>fine i</sub> = the mass of the ne earth per volume of ne earth of the depth increment i  
(g ne earth cm<sup>-3</sup> ne earth = dry soil mass [g] – coarse mineral fragment mass [g]) / (soil sample volume [cm<sup>3</sup>] – coarse mineral fragment volume [cm<sup>3</sup>])

vGi = the volumetric coarse fragment content of the depth increment i

t<sub>i</sub> = thickness (depth, in cm), of the depth increment i

= conversion factor for converting mg C cm<sup>2</sup> to Mg C/ha.

When SOC stock changes are monitored, changes in bulk density should also be considered. Because of bulk density variation, comparison of SOC stocks should be made on an equivalent soil mass basis (ESM). This means that SOC stocks over time are compared to the same mass of soil. This method fixes the effect of SOC content and bulk density variation in different soil depths (Wend and Hauser 2013). Overall, SOC stock determination should always be conducted with the same sampling and analyzing protocol, so that changes could be detected reliable (FAO 2019).

### 6.1.3 Minimum number of sampling points

A long monitoring period and a large sample size are needed for evaluating soil treatment effects on SOC (due to large spatial variation of SOC). The smallest difference in SOC stock that can be detected and is statistically significant, is based on the minimum detectable difference (MDD), and it can be determined through power analysis (equation 2) (Zar 1999).

$$MDD \geq \frac{s}{\sqrt{n}} \times (t_{\alpha, v} + t_{\beta, v}) \quad (2)$$

where

MDD = minimum detectable difference

S = standard deviation of the difference in SOC stocks between t<sub>0</sub> and t<sub>1</sub>

n = number of replicates

v = n – 1 is the degrees of freedom for the relevant t-distribution

t = values of the t-distribution given a certain power level (1-β) and α level.

The minimum number of samples needed to detect the difference of two different sampling points can be calculated with equation 3 (Vanguelova et al. 2016).

$$n \geq \left( \frac{S \times (t_{\alpha} + t_{\beta})}{MDD} \right)^2 \quad (3)$$

where

n = number of samples,

MDD = minimum detectable difference

S = estimated standard deviation,

$t_{\alpha}$  = two-sided critical value of the t-distribution at a given significance level ( $\alpha$ ) frequently taken as 0.05 (5%)

$t_{\beta}$  = one-sided quartile of the t-distribution corresponding to a probability of type II error  $\beta$  (e.g. 90%).

There are different calculation methods, and for example Mäkipää et al. (2008) used the equation 4 to calculate the number of plots needed for detection of soil carbon stock changes in Finnish forests.

$$n = (t \times s/E)^2 \quad (4)$$

where

n = number of plots required

t = value from Student's t distribution table (number of degrees of freedom and confidence interval considered)

s = estimated standard deviation

E = desired half of the confidence interval.

Schrumpf et al. (2011) took soil samples from 12 sites from CarboEurope Integrated Program across Europe to find out sufficient core number in plot/field scale and if the equivalent soil mass method would increase the smallest detectable change. They took 100 sampling points per site (up to 60 cm depth) and they covered the major land use types, deciduous and coniferous forests, grasslands and croplands. The authors concluded that in cropland sites the spatial variability was smallest which also led to the lowest minimum detectable difference ( $105 \pm 28 \text{ g C/m}^2$ ). In grasslands the minimum detectable change was  $206 \pm 64 \text{ g C/m}^2$  and forest sites  $246 \pm 64 \text{ g C/m}^2$ .

## 6.1.4 Sampling frequency

A single re-sampling would not distinguish any interannual variability and long-term trends. Repeated soil inventories during a certain period would be advisable, instead of just one re-sampling after several years. More frequent sampling will increase precision (Schrumpf et al. 2011). SOC change between two sampling points should be greater than MDD, which means that if the expected stock change rate is low, the sampling time is also less frequent. Carbon input, climate and seasonal weather influence on the carbon gained or lost. Under variable environmental conditions, a longer sampling frequency is needed, to ensure that any changes in SOC stocks can be detected (FAO 2019). Intra-annual variation is also important to consider. SOC decomposition is mainly moisture and temperature dependent (Paul 2007), farming practices, carbon inputs, and carbon inputs due to natural reasons vary seasonally. Repeated sampling over several years should be planned to minimize intra-annual variation; this can be done by ensuring that sampling is conducted during the same season or by comparing more than two years (van Wesemae et al. 2010).

Smith (2004) studied the increased carbon input effects on the SOC stock change detection between two sampling points, and reported that when the C input increased by a maximum of 20 to 25%, changes in SOC stocks could be detected after 6-10 years (with 90% confidence) (Smith 2004). Schrumpf et al. (2011) also found out in their vast soil monitoring study (mentioned above) that general trends in soil organic carbon indicate that with 100 samples (per site), the change would be detectable after 2-15 years (10 cm depth). In a depth of 30 cm, the time varied between 7 years (grasslands) and 14 years (croplands, conventional farming methodologies) to 20 years in forests (Schrumpf et al. 2011).

## 6.1.5 Uncertainties in soil sampling

The absolute true mean value of SOC stock is not possible to determine and reporting the average value and measure of uncertainty is recommended (FAO 2019). Soil sampling includes several steps and each step includes possible error sources. Table 1. summarises the systematic error sources at different scales. Identifying different potential sources of uncertainties is important when considering the possibilities to reduce them. Biggest problem based on literature can potentially be to neglect bulk density determination. Adequate sampling depth is also under debate (Vanguelova et al. 2016).

**Table 1.** Possible soil sampling error sources at different scales. Modified from Vanguelova et al. 2016.

<b>Sample</b>	<ul style="list-style-type: none"> <li>Bulk density is not assessed</li> <li>Samples are not homogenized</li> <li>Different analytical procedures</li> <li>Coarse fragment volume is not assessed</li> <li>Different soil horizons and layers are not separated accurately</li> <li>Inappropriate time for soil sampling</li> </ul>
<b>Profile</b>	<ul style="list-style-type: none"> <li>Sampling at not full soil depth</li> <li>Sampling by horizon versus soil depth</li> </ul>
<b>Plot</b>	<ul style="list-style-type: none"> <li>Bulk density and stone content not analyzed</li> <li>Not enough sampling points</li> <li>Different sampling schemes</li> <li>Small scale variability not accounted</li> <li>Measurement error including sample preparation</li> <li>Not harmonized inventory teams</li> </ul>
<b>Landscape/National/ European</b>	<ul style="list-style-type: none"> <li>Lack of local and regional representativeness of sampling plots</li> <li>Important areas are underrepresented (e.g. peat soil)</li> <li>Lack of forest cover maps and accurate soil/hydrology maps</li> </ul>

An appropriate scale related to carbon stock monitoring is important to define. Different scales could include profile, plot, forests, catchment, national or wider areas. Soil heterogeneity and spatial variability are important to take into consideration, as the factors are clearly scale dependent (Goidts et al. 2009). In general, soil properties vary more with increasing study area. According to Hobley and Willgoose (2010), soil carbon’s spatial variability can rise sevenfold when scaling up from a point sample to landscape scale. This can lead to high uncertainties in SOC stock calculations if scaling up is not considered accordingly (Vanguelova et al. 2016).

Davis et al. (2018) compared soil organic carbon measurement protocols in U.S. and Brazil and they found out that reported procedures reflected big variabilities, which makes it hard to compare results from different study sites. Differences may be due to different sampling protocols instead of differences in soil carbon stocks.

## 6.1.6 Sampling cost

Mäkipää et al. (2008) calculated the sampling costs for plot scale soil sampling in Finnish forests. Costs of soil sampling depends on several factors and those costs can be divided into fixed and variable components. Fixed components include costs that are not dependent on the number of the soil samples ( $n$ ). For example, direct personnel costs from transportation, accommodation and driving time to the sampling site (salary). Variable costs are dependent on the sample size ( $n$ ) and it includes the costs of soil sampling, sample preparation and analyses in the laboratory (salary and all other costs). According to Mäkipää et al. (2008) measuring the carbon in soil organic layer costs 520 €/plot with 10 analyzed samples. In this study and with their sampling protocol the minimum detectable change was  $>860 \text{ g C/m}^2$ , which is not very good precision. If sample size per plot was increased to 30 sampling costs arise to 1100 € and detectable change drop to  $540 \text{ g C/m}^2$ . In Peltoniemi et al. (2004) study, the organic layer increased in average by  $47 \text{ g C/m}^2$  during a 10-year period and this relatively slow change makes it hard and costly to detect on a small scale.

Singh et al. (2012) calculated how much would cost to measure field level soil carbon stock size in cropping field in Australia. They concluded that sampling to 30 cm depth in a 68-hectare area with  $< 2 \text{ t/ha}$  standard error, the cost would be 2500 AU\$ (~1500 €).

## 6.2. Biomass carbon stock estimation

The conventional methods to estimate amount of biomass are harvesting to determine exact biomass and estimation approaches which rely sampling and statistics. Harvesting and directly measuring all biomass (weight and volume) is destructive method and it gives accurate values (Husch et al. 2003). This type of approach is laborious and time consuming, but it provides basic data information to the different applications. For example, different allometric equations and growth models are conventionally used methods for biomass estimations and they are developed from information provided by destructive methods) (Mäkelä and Valentine 2019). Destructive methods are part of the basic research and needed for better model development, model validation and to increase our knowledge of vegetation characteristics etc., but they are not suitable for biomass estimations in a purpose to estimate permanent carbon pools. Other forest field method, that don't include harvesting, is measuring diameter and height of trees and to identify different species and then link those measures to other three attributes like, total biomass, via allometry (Husch et al. 2003).



Husch et al. (2003) pointed out that the most economical approach to estimate aboveground forest biomass is to use data from forest inventories. This inventory-based approach would increase the usage of already gathered information and information gathered for other purposes. National forest inventories are conducted in several countries and they have been common practice in many places for decades (Tomppo et al. 2010). On the other hand, several countries where carbon sequestration potential is high, forest inventories are not a common practice, which hampers the application. Small scale forest inventories are also conducted to research purposes (Tomppo et al. 2010).

## 6.2.1 National forest inventories

National forest inventories (NFI) are conducted to provide information about forest resources. Most basic variables that forest inventories produce are related to forest area and growing stock volume. Inventories are based on large field measurements, where tree parameters are measured. Field plot measurements and systematic or random sampling are ways to produce information which can be upscaled to comprehend large areas. In national forest inventories, the whole country is covered with regular networks of plot measurement clusters. These field measurements can be used for reliable forest statistics and calculations for larger areas (Tomppo 2014).

Information from forest inventories are usually used for example policy making, forest management planning, assessing sustainable forestry, greenhouse gas and carbon stock evaluation and research (Tomppo 2014). The first NFI in Finland was carried out in 1921-1924 (Tomppo et al. 2010). Current day inventory in Finland is multisource inventory where several data sources are utilized (field measurements, satellite data and digital maps) (Tomppo 2014). Most European countries conduct forest inventories, for example Austria (first 1952-1956), Sweden (1923-1929), Spain (1965-1974), Great Britain (1924) and Italy (1986-1988). USA (1928) and Brazil (1980s) are also managing their forest resources through inventories. Canada is one of the biggest forest countries, where national forest inventory is not mandated through legislation. China (first 1973-1976), Japan (1951), The Republic of Korea (1960s), New Zealand (1946-1955) and Russian Federation (2007) are countries in Asian continent that conduct inventories (Tomppo et al. 2010).

## 6.2.2 Allometric equations

Allometric equations are mathematical models that describe the relationship between tree characteristics that are easier to measure to another tree properties that are hard to

measure. This relationship is typically based on detailed measurements of vegetation, where small sample size is representing a population of interest (Mäkelä and Valentine 2019). Easier tree measurements are for example diameter and total tree height and harder ones are volume or biomass. Diameter is typically measured from chest height and the total height can be estimated with a hypsometer and leveling rod. For example, biomass equations are developed from harvested and weighted vegetation samples. Each sample is oven dried, weighted with high detail, stem, stump, roots, branches and foliage separately, and after sufficient number of samples, some consistency can be seen. With regression techniques, certain parameters of allometric equation that relate biomass and measured variables can be found (Birdsey et al. 2013). Each allometric equation is as good as its parameters are (Mäkelä and Valentine 2019). Individual tree estimates can be expanded to larger areas by knowing the probability of sampling each tree (Birdsey et al. 2013).

Different biomass or volume equations are a cost-effective way to evaluate large areas, but there is scarcity of representative equations. Population of trees under interest maybe different than population from where the equation was developed, and if only few equations are available, there is a problem. This issue is particularly true in tropical regions (Rex et al. 2020). Generalized biomass equations can be used when local or species-specific biomass equations are not available (Birdsey et al. 2013). Biomass equations are typically presented to individual species, groups of species (Pearson et al. 2007) or for geographic regions (Duncanson et al. 2015). Allometric equation are also typically developed to forest trees which diameter at breast height (DBH) is bigger than 10 cm and equations for smaller trees are rarely available. This means that forest understory vegetation is difficult to estimate cost-effectively (Han and Park 2020).

One example of allometric equation which can be used to calculate oven-dry tree biomass  $M$  (kg) according to Brown (1997) is shown in equation 5.

$$M = a + bDBH + cDBH^2 \quad (5)$$

where

DBH = diameter (cm) at breast height (1.3m)

a,b,c = best fit parameters.

Biomass growth can be also simulated with growth models. Growth models combine carbon allocation models to factors affecting carbon accumulation (Mäkelä and Valentine 2019).

### 6.2.3 Estimating carbon stock from forest inventories

Most forest inventories are focused on timber estimation, but constant need for information about forest health, soils, wildlife and other nontimber values have created development of integrated or multisource inventories. Information from a timber inventory is insufficient for a complete estimate of a carbon stock, because they usually estimate only the volume of the main stems, ignoring other components of the vegetation and other carbon pools in the ecosystem. Carbon stock estimations need an inventory of the total biomass of live standing timber, biomass of the understory vegetation and estimations of dead biomass, root biomass and soil carbon pools. Total biomass can be adjusted with an expansion factor to include all other biomass quantities (Husch et al. 2003) or other carbon pools could be also estimated by conducting additional measurements at the inventory (Birdsey et al. 2013). If inventories are conducted so that other biomass is ignored, to include weights of these components (roots, foliage, understory vegetation and detritus on the forest floor) the weight of the commercial volume should be multiplied by an expansion factor which typically varies from 1.3 to 2.5 depending on species, forest age, average tree height and amount of dead matter. Expansion factors are generated from allometric relations (Husch et al. 2003).

Carbon pools locates in vegetation biomass aboveground and belowground, and each pool needs different sampling methods. Size of those carbon pools vary, and it should be decided if the certain pool is cost effective to estimate for certain purposes. Live trees and their roots are important to include to all activities. Understory vegetation and nontree biomass maybe be beneficial to measure and monitor only if they are significant component of the total biomass. This is the case in areas where the main biomass consists mainly from shrubs and other form of nontree vegetation. Some forests may mainly composite from big trees and understory vegetation is not a large part of total biomass. The forest floor should be included in carbon pool accounting in most cases, especially in conifer dominated forest, because it is known that in this type of forest the biomass in forest floor consists big part of the total pool. Understory vegetation in the forest inventories can be measured with harvesting technique, where small subplots are harvested, and vegetation is oven dried, pooled to composite sample and weighted. After measuring, the information can then be upscaled to the whole plot (Pearson et al. 2007).

Amount of biomass can be calculated with equation (8) and multiplying by 0.5 the metric t/ha for the amount of carbon (Pearson et al. 2007).

Oven-dry weight (g) of biomass / sampling frame area (cm<sup>2</sup>) \* 100 (6)

where

multiplying by 100 converts the unit to metric t/ha.

Below-ground biomass comprehends coarse and fine roots and they are included to carbon pool accounting by applying a regression models which links belowground biomass to aboveground biomass. Example Cairns et al. (1997) developed regression models that can be used in different forest biomes (7-9):

$$\text{Boreal: BGB} = \exp(-1.0587 + 0.8836 * \ln\text{AGB} + 0.1874) \quad (7)$$

$$\text{Temperate: BGB} = \exp(-1.0587 + 0.8836 * \ln\text{AGB} + 0.2840) \quad (8)$$

$$\text{Tropical: BGB} = \exp(-1.0587 + 0.8836 * \ln\text{AGB}) \quad (9)$$

where

BGB =belowground biomass density in t/ha

AGB= aboveground biomass density t/ha

(n=151; R<sup>2</sup>=0.84.).

Dead biomass comprehends dead organic matter in forest floor and dead trees (on the ground or standing). Forest floor dead biomass should be estimated like living forest floor biomass. Most time-efficient way to estimate carbon stocks in dead wood is the line intersect method, where each dead wood intersecting at least 100 m length (per plot) line is measured and classified via density (Harmon and Sexton 1996). There are different density classes and they are developed by forest scientists and they are based on different decomposition models of dead wood (Beets et al. 1999).

Most accurate and precise carbon stock estimation of trees is achieved with direct methods where all trees (above a minimum diameter) in sample plot are measured. The minimum diameter varies according to trees that are expected to be found in a sampling area. Environments where the trees grow slower (e.g. arid) the minimum diameter may be 2.5 cm and in humid environments, where the tree growth is fast, the value may be up to 10 cm (Pearson et al. 2007). Tree biomass is often estimated with equation where only the diameter at chest height is used as a variable. Height and diameter as the independent variables result better estimates but measuring the tree height increases the cost of monitoring. If there is vast monitoring network with plenty of data, the regression equation with diameter only can result a high significance (Pearson et al. 2007), but if not the variation of total biomass estimations could be high. For example, the total estimate of tropical forest carbon biomass stock varies by 35.3 Pg depending if the height is included (Feldpausch et al. 2012).

## 6.2.4 Inventory planning and sample size

Inventories are conducted in a purpose to provide information and that needed information defines how it is gathered. Available funds and the costs of an inventory will influence the

chosen design. Main factors affecting the costs are precision, total size of the area and the minimum size of the unit area where estimates are required (Husch 2003). The accuracy requirement defines the number on needed sample plots. Keller et al. (2001) calculated the required number of plots and total sampling area for biomass estimations if total error is within 20% of mean with 95% confidence (table 2). When the area increases the total number of required samples decreases.

**Table 2.** Required number of sample plots and total area to meet error less than 20% (Keller et al. 2001).

Plot size (ha)	n	Total area (ha)
0.09	43	3.87
0.25	21	5.25
0.49	15	7.35
1.00	10	10.00
1.96	8	15.68
4.00	6	24.00

## 6.2.5 Uncertainties in forest inventories

Typical errors sources in forest inventories are sampling error, measurement error and prediction error from used models. Classification error of remote sensing imagery is also one error source, if remote sensing is used. The essential problem in inventory-based approach is that obtained samples should represent the population. If samplings are representative, useful statements can be made about characteristics of the population, like volume or weight per unit area, number of trees etc. These characteristics, parameters, exact values would be known if the entire population would be measured, but due the time and cost factors, sampling provides estimated values for these parameters. Estimates are calculated from samples and these statistics are summary values which represent the whole population. If determined parameters are not representative it will lead to a sampling error. That's why the efficient sampling design is important. The used sampling units, the number of samplings, the manner of selecting and distributing the sampling points over forest area, measurement and result analyzing procedures are all important parts trying to decrease the sampling error (Husch 2003).

Keller et al. (2001) calculated the size of the sampling error and other error sources (table 3) in the Amazon area field measurements and data analysis. Their results show that in mean

sampling error was approximately 15%. Field plot aboveground estimation's uncertainties has been reported to be even 20-30% (Keller et al. 2001, Chave et al. 2004).

Table 3. Biomass density estimations with related errors.

	<b>Mass (Mg/ha)</b>	<b>Sampling 95% CI (Mg/ha)</b>	<b>Other error (%)</b>
Trees (DBH≥35cm)	177	24	20
Trees (15<DBH<35cm)	47	10	50
Trees (DBH<15cm)	40	8	50
Vines and epiphytes	18	2	50
Dead fine AGB	8	2	50
Dead coarse AGB	19	3	50
All below-ground	63	9	50
<b>Total biomass</b>	<b>372</b>	<b>56 (15%)</b>	

Measuring tree height is time consuming and it's possible that height is not measured form all trees form inventory plots (Sullivan et al. 2018). For example, according to The Amazon Forest Inventory Network RAINFOR, the guideline is to measure the height of 40 trees in 1 ha area if time prevents all trees being measured (Phillips et al. 2009) which means that in tropical forests 90% of tree heights are not measured but predicted. Different prediction models perform differently, and the performance can be estimated by calculating prediction error. For example, root mean square error (RMSE) can be used to describe the difference between measured and predicted heights (Sullivan et al. 2018).

Measurement errors include errors that arise from defects in the sampling procedure, like mistakes in data collection or processing. Measurement errors don't decrease when increasing the sample size. For example, appropriate training of field crew, quality control, appropriate mathematical models and well-prepared maps would reduce the nonsampling errors (Husch 2003).

## 6.2.6 Accuracy and cost

In-situ sampling costs depend on design, the number of attributes to be collected, salary levels and on the accessibility of the forest area. Costs increase with accuracy. Berenguer et al. (2015) studied how forest carbon stock could be quantified through field measurements in cost-effective way. They conducted the study in Amazon, which is area lacking intensive field measurements. Several tropical countries are suffering this lack of filed measurement. Study site included 224 sampling plots spread evenly to two 5000 ha areas (three different

forest type). They conducted field measurements to all biomass carbon stocks and soil. These results were compared to simulations where single value for wood density was used, without identifying the stems. They also compared the field measurements to default values, defined by FAO and IPCC. The total cost quantifying carbon stocks of field sampling by assessing each forest component was 364 000 US\$ (~311 000 €) with 224 0.25 ha forest plots. They calculated that without species identification the costs would decrease by 58%. This reduction would be due to decrease in salary costs when experienced taxonomic experts are not needed. They also noticed that forest soil was by far the most expensive and time-consuming part to measure (with conventional soil samples and dry combustion method). In their study, the soil sampling cost approximately 2250 US\$ (~2680 €)/ha and identification and measuring large and small stems cost ~500 US\$ (~595 €)/ha each.

Berenguer et al. (2015) also expressed the average error compared to intensive field measurements with unit Mg C/ha. Field sampling with species identification gave errors close to zero (accurate values were not presented in article). Protocol where stems were measured but not identified gave average errors of 2.69 Mg C/ha, 6.42 Mg C/ha and 14.22 Mg C/ha which represent 3%, 5% and 31% of the total carbon stocks contained in those stems (average from different areas, three different forest types). FAO default values gave average errors of 21.16 Mg C/ha and 5.02 Mg C/ha (value for two different areas, forest type doesn't matter in default values). IPCC values performed poorly, average errors were 50.16 Mg C/ha and 34.02 Mg C/ha.

## 7. Modelling soil carbon stocks

Soils are very heterogeneous, and it would take a large number of direct measurement samples to estimate the size of the soil carbon pool reliably. Changes in soil carbon pools occur slowly and monitoring it is difficult (Mäkipää et al. 2004). Modelling soil dynamics and simulating stock changes helps to tackle those obstacles, and offers a mathematical way to estimate SOC, but completely accurate model is difficult to develop, because the large uncertainties in empirical data and the complexity of the carbon turnover process in soil (Peltoniemi et al. 2007).

### 7.1. Three levels of soil models

FAO (2019) categorizes soil organic matter models into three levels according to different models of approach. These categories are 1. empirical models, 2. soil process models and 3.



ecosystem models. Model categorizing varies, and for example Paustian et al. (2019) divides models that predict SOC changes into empirical models and process-based models.

Empirical models represent the observed relationship between carbon stocks and environments. Empirical models are based on statistical relationships and describe how environmental and management variables interact with SOC stocks and SOC stock changes (FAO 2019, Paustian et al. 2019). Statistical relationships are estimated from field experiments, where SOC changes due to environmental and management variables are observed (Paustian et al. 2019). Temperature, precipitation, soil clay content and land use are typical studied variables affecting SOC changes and empirical model development (FAO 2019). Best known empirical approach is the model developed by Intergovernmental Panel on Climate Change (IPCC) to estimate SOC stock changes for the national greenhouse inventories (IPCC 2003). Empirical type of approach has been the basis for more complex models (FAO 2019). The drawback of using empirical models is that typically these equations are generated for specific soil types, climates, management and carbon inputs which leads to the situation that models are not working adequately when those variables change (FAO 2019). Other limitation is lack of field experiment data from many climates, soil types and management combinations (Paustian et al. 2019). Empirical models developed by IPCC (2003) are developed from global data sets, and they are mentioned for national scale application. Thus, regional or local scale SOC stock estimation need new estimations of parameters used in models (Paustian et al. 2019).


Process-based models are models that estimate the SOC stock changes aided by SOC dynamics through time. These SOC dynamics consider the effects of climatic and soil factors with land use and management variables (Paustian et al. 2019, Senapati et al. 2014). These models are more detailed, and they determine SOC stocks and changes by using mathematical functions (sets of different equations), where physical and chemical soil processes are considered (FAO 2019). These models are usually constructed from several compartments (Fig. 7) where each represents the fraction of SOC with similar characteristics. Decomposition rate and mechanism driving carbon stabilization are typical factors directing model compartment division (Stockmann et al. 2013). The carbon flow in models moves from litter to the microbial pool and then to soil carbon pool. Moving from first compartment to next, the stability of soil organic carbon increases (FAO 2019). Yasso (Liski et al. 2005) and RothC (Coleman and Jenkins 1996) are examples of process-based models (table 4). FAO recommends that these types of models should be used when required (model specific) data is available.

Most models are developed for research purposes where aim was to study how different changing variables affect SOC dynamics. Under interest has been how SOM functions with environmental variables, e.g. temperature and moisture, edaphic variables. e.g. pH and soil texture, and land-use and management practise, e.g. vegetation type, productivity, tillage



and residue management. Process models integrate these factors and also controls affecting decomposition and organic matter stabilization in soils into one model (Paustian et al. 2019). Even though dynamic process-oriented models might be quite comprehensive, they don't include all important ecological processes affecting soil carbon. Some exclusions might include e.g. biomass growth and nutrient cycle (FAO 2019). Process oriented models also need relatively few data requirements. Information about climate, soil and productivity are typical data requirements for simulations (FAO 2019).

Process based models can be further developed by integrating models with several data sources (Campbell and Paustian 2015) like flux measurement networks and existing long-term field experiment (Harden et al. 2018)



**Figure 7.** Separation of the SOM in different compartments representing different decomposition rates. Modified from Willgoose 2018.

Ecosystem models present the third level of modelling according to FAO (2019). Ecosystem models simulate stock changes in time, considering the same factors as the level two models, but they also integrate above- and belowground plant biomass growth and carbon inputs, water and nutrient dynamics and their interactions. CENTURY (Parton et al. 1987) is one of the existing ecosystem models used for SOC estimations. Models with several compartments, processes and interactions to simulate, need higher amounts of soil, climatic

and management data. Higher complexity can provide higher accuracy, but great data requirements can be difficult to obtain, which limits the application (FAO 2019).

More than 250 models describing soil carbon turnover (Manzoni and Porporato 2009) have been formulated, each for different purposes. Models differ from each other based on the biochemical and physical processes and the underlying assumptions. Table 4 summarises examples of commonly used named models. Yasso (Liski et al. 2005) and RothC (Coleman and Jenkins 1996) are process based models, CENTURY (Parton et al. 1987) is ecosystem model and FullCAM (Richards et al. 2004) is combination model which integrates several different models into one full ecosystem model.

**Table 4.** Overview of different types of process-oriented models. Yasso and RothC are process models, CENTURY is ecosystem model and FullCAM is combination of several models.

	Model name			
	Yasso	RothC	CENTURY	FullCAM
<b>Short description</b>	Dynamic soil carbon model. Simulates decomposition of organic matter.	Models the turnover of organic carbon	Ecosystem model. Simulates different plant-soil systems C, N, P and S dynamics in a long term.	Ecosystem full carbon model. An integration of biomass, decomposition, soil carbon models and accounting tools.
<b>Land use area</b>	Forest	Crop, forest, grassland	Grassland, forest, crop, savanna	Forest, crop, agroforestry
<b>Input variables</b>	1) Soil C stock beginning, 2) C input to soil, 3) climate	1) monthly rainfall, 2) monthly open pan evaporation, 3) average monthly mean temperature, 4) clay content of the soil, 5) an estimate of the decomposability of the incoming plant material (the DPM/RPM ratio), 6) soil cover, 7) monthly input of plant residues, 8) monthly input of farmyard manure and 9) depth of the soil layer sampled.	1) air temperature (monthly average maximum and minimum), 2) monthly precipitation, 3) lignin content of plant material, 4) plant N, P, and S content, 5) soil texture, 6) atmospheric and soil N inputs and 7) initial soil C, N, P and S levels and 8) litter or crop residues	All that: forest physiological growth model 3PG, the carbon counting model for forests CAMFor, cropping and grazing systems CAMAg, the microbial decomposition model GENDEC and the Rothamsted Soil Carbon Model RothC require.
<b>Time step</b>	Year	Month	Month	?
<b>Depth</b>	1 m	0-20 cm (topsoil)	0-20 cm (topsoil)	0-20 cm (topsoil)
<b>Ref.</b>	Liski et al. 2005	Coleman & Jenkins 1996	Parton et al. 1987	Richards 2004

## 7.2. Soil carbon turnover models internal structure

According to Batlle-Aguilar et al. (2011) SOC turnover models can be divided into categories based on their internal structure. Processes-oriented (multi-compartment) (Smith et al. 1998), organism-oriented (Post et al. 2007), cohort (describes decomposition as a continuum) and a combination of the first two, are four categories that each describe SOM dynamics models.

Process-oriented, or compartment models, are built to consider SOM transformation and migration through different soil layers. A compartment refers to different fractions of SOM that each have different chemical and physical characteristics. These kinds of models can be complex, including multiple compartments, or a simple one compartment, or even no compartment models, where degradation is assumed to be a continuum (Smith et al. 1998). Process-based models can be combined with Geographical Information Systems (GIS), which adds benefits for regional-scale studies. CENTURY (Parton et al. 1987) and RothC (Coleman & Jenkins 1996) are process-based models combined successfully with GIS (Batlle-Aguilar et al. 2011). As a downside, theoretical compartments created to describe SOM dynamics are difficult to compare with actual measurements of soil fractions. Thus, validation and testing are quite limited leading to a situation that model can include an undefined inaccuracy (Batlle-Aguilar et al. 2011).

Organism-based models describe SOM movements between pool, and each of these organism pools is classified. The main drivers (soil fauna) of SOM fluxes and transformation are explicitly accounted for, which gives more accurate estimations where the model is based on (Post et al. (2007). Accuracy is the main advantage and the negative side is that the relationship between the abundance of soil biota and degradation rate of organic matter is not yet commonly agreed upon. Another downside in organism-oriented models is that site-specific calibration requires the characterization of the whole soil fauna with complex techniques. Compared to process-based models, the relationship between degradation rate and substrate concentration is well known. First-order kinetic rate (assumption: the bigger the concentration, the bigger the decomposition rate) (Senapati et al. 2014) is simple and often a suitable way to describe the organic matter transformation. Rate of these reactions can be determined in a laboratory (e.g. litter bag) and then used in process-based models, but organism-oriented models need complex site-specific calibration. Process-oriented models are more popular because of their usability (Batlle-Aguilar et al. 2011).

Cohort models divide the organic matter in soils into cohorts, and further into different pools (Senapati et al. 2014). In a cohort model, the microbial physiology is considered to be the

main driving factor in organic matter decay. The fourth group of models is a combination of the process- and organism-oriented models, which have high specific data requirements and thus not commonly used (Batlle-Aguilar et al. 2011).

In most soil carbon models the size of the microbial pool (microbes, fungi and fauna) is not determined, but the decomposition is considered by variables that affect the microbial activity (e.g. temperature and soil moisture). This means that the microbial activity in model is expressed as decomposition rate of compartment (Peltoniemi et al. 2007). Because SOM is complex and includes fractions of different stability, the energy needed for decomposition varies also (Davidson & Janssens 2006, Peltoniemi et al. 2007). In model compartment development, these different decomposition rates and needed energy requirements are approximated and then divided to several different compartments that differ in organic matter turnover time. Because of the complexity of SOM, several studies confirm that simulation of carbon dynamics cannot be adequately approached with one compartment only (Kätterer et al. 1998, Davidson and Janssens 2006), but more complex SOM models, with several pools, are necessary when modelling carbon concentration changes in soils and the atmosphere (Schimel et al. 1994). Large soil carbon stocks located in slow turnover pools in soils and fluxes represent the fast turnover carbon pools (Fig. 7). Models with one pool and one turnover rate will overestimate the carbon response because changes in the stable stocks happen in the slow pool (Telles et al. 2003)

## 7.3. SOC model scales

SOC models can be formulated to different scales. According to Campbell and Paustian (2015) three commonly used scales are microsite, ecosystem and global. Each scale has its own limitations and use. Microsite is the smallest scale and models designed for that can be used, for example, to predict short-term and small changes. Microsite presents, for instance, a small area of a rhizosphere. These small-scale models are difficult to link into larger scale dynamics and microsite models are also dependent on specific soil fractionation method (a method where different soil organic matter particles are fractionated from each other).

Ecosystem scale models can be used to model a hypothesis based on mechanistic or empiric relationships and predict impacts of changes specific to a certain site. Ecosystem scale limitations are required for site-level data, and on this scale, models cannot represent mechanistic relationships that are important in smaller scales. RothC (Coleman and Jenkins 1996) and Yasso (Liski et al. 2005) are examples of ecosystem scale SOM models. SOM models can also be formulated on a global scale. Global scale SOM models that can be used to model

hypotheses of large-scale dynamics, simulate global scenarios and predict climate change with dynamic soil feedback (Campbell and Paustian 2015).

The scaling decision is a critical component of the simulation of soil carbon for inventory purposes (Peltoniemi et al. 2007). Input data and model application will determine to which scale the soil carbon stocks can be estimated. Many models are developed to cover a small spatial resolution and a short time period (daily time step), but those models can be scaled to a larger resolution. Scaling-up can create biases, Ogle et al. (2006) concluded that increasing model spatial resolution with coarser-scale parameters can lead to significant biases. Rescaling is possible but should be done with careful consideration of the uncertainties (Peltoniemi et al. 2007).

## 7.4. Model selection

The choice of modelling approach depends on the purpose, available resources and expertise. FAO (2019) recommends that a locally validated model should be preferred. Internal model calibration, with region(site)-specific data, and factors adapted to that give more accurate results (FAO 2019).

Models need several types of data and data availability is an important factor when deciding which approach to use (FAO 2019). Several different process-based models could be used for soil carbon inventories, but the selection is constrained by the availability of model input and evaluation data (Peltoniemi et al. 2007). Data sources can be, for example, national forest inventories (NFI), soil samplings and remotely sensed data. There are several international soil and climate databases that provide data for model inputs (e.g. FAO Global Soil Information Carbon Map, Solid Grids- Global Soil Data Facility). These databases do not provide data for all situations, but in some cases local data may be available ((Peltoniemi et al. 2007). National forest inventories are also widely conducted in several countries (Tomppo 2010).

Typically soil carbon models need input data, parameters and test data (FAO 2019.). Input data is data that a model needs to output predictions. Most typical input variables are temperature, moisture, soil texture and nitrogen. Those are the main factors affecting the decomposition processes (Peltoniemi et al. 2007). Information about land management and disturbances are also key model inputs, because minor changes in land-use may lead to major changes in soil carbon (Peltoniemi et al. 2007). Thus, forest inventories with land use surveys, remotely sensed data and long-term soil surveys are important data sources. Data from several sources also helps up-scale models. Amount and coverage of biomass, species composition, topographic position, temperature and thermal regimes and edaphic

characteristics are important factors and data sources that provide information on different scales are needed when plot level data is scaled over large areas. Long-term experiments give valuable information and measured data can be used to evaluate and compare models. Measurements are also needed for model development and more extensive data gives better building blocks for new models. SOC models are usually used in larger, ecosystem scale, carbon change simulations and future predictions (as it is possible to formulate site-specific calibration, see above) (Peltoniemi et al. 2007). Parametrization is a critical step in the model development. If parameters are not sufficient, new values might need to be measured or parameters to be fitted with existing calibration data (FAO 2019). Test data is the data that the model has predicted (e.g. SOC stock and changes) and it can be divided into calibration data and validation data. Test data is used to test model outputs (FAO 2019).


Most commonly used SOC models are developed for temperate climate, which means that they are parametrized and calibrated with data that fits in certain conditions. For tropical and subtropical applications, these models should be validated for conditions corresponding area under interest (Campbell and Paustian 2015).

## 7.5. Uncertainties and sensitivity analysis

There are several possible sources of uncertainties in SOC models. Some errors in SOC models are not avoidable because they arise from problem that man can't exactly describe complex chemical, biochemical, physiological and biological systems in a mathematical way. In general, there are two main sources of uncertainty: model uncertainty (mentioned above) and uncertainty of modelled system inputs (FAO 2019). Model structure includes uncertainties because of real life processes can't be thoroughly represented in models. Model uncertainties also includes parameter value uncertainty, which means that the correct value of the parameter that determine the model estimations is imprecise. Modelled system inputs include measurement errors and natural variability (Ogle et al. 2010). According to (FAO 2019) structural uncertainty is the one which causes major difficulties. That's because if the processes are not adequately represented, no effort will not reduce the uncertainties. Structural uncertainties can't be ignored because if models are used in carbon storage change estimations in a purpose to mitigate climate changes, it is important to recognize uncertainties and adjust buffers that are big enough to secure the negative result in carbon budget.

Sensitivity analysis (Fig. 8) is a useful tool to identify the most significant variables and parameters for further analysis. FAO (2019) recommends that model sensitivity analysis and uncertainty assessment is conducted for every simulation scenario to confirm that model is

suitable for its application. Information about model inputs and processes is also important to evaluate continuously.



**Figure 8.** Illustrative figure of model (RothC) uncertainty and sensitivity presented with typical parameters needed to simulate carbon turnover. In this figure, plant carbon input is a parameter that has high uncertainty and high model sensitivity, which makes it problematic. Modified from FAO 2019.

Models should be calibrated to reduce uncertainties. Calibration can be conducted by calibrating model parameters that are most effective for the model performance (FAO 2019). Site specific calibration means that the simulation of SOC dynamics in a certain area requires calibration with data specific to that soil area, like soil particle size distribution, pH, soil type, decomposition rate, etc. This kind of information input makes models more reliable, but it complicates the prediction of changes. To truly predict SOC changes, a model should be able to simulate dynamics without site-specific data input (Smith et al. 1997). If models that need site-specific calibration are used in long term simulations, it may cause major biases in the results if the site changes over time (Willagoose 2018).

## 7.6. Model performance

SOC model's performance can be studied by comparing values from simulations to actual measured values from long-term field experiments. Commonly used measures of model



performance are the coefficient of determination ( $R^2$ ) and root mean square error (RMSE) (122.). Smith et al. (1997) compared the performance of nine different SOM models. A model's performance was evaluated with different data sets (arable land, forest soil, strong fertilization, etc.) and the results were compared to soil samples (from long-term soil inventories). All nine were process-oriented multicompartiment models. They concluded that their tested models could be divided into two groups according to their performance. SOMM, ITE and Verbene were the most poorly performing SOM models and RothC, CANDY, DNDC, CENTURY, DAISY and NCSOIL showed more accurate results. From those nine models, RothC and CENTURY are the most interesting ones considering carbon dynamics. RothC, CENTURY and CANDY did not show significant biases in overall simulations and they performed the best in all datasets (Smith et al. 1997). RothC and CENTURY can be used in different land-use areas, which explains partly their good performance. One source of error occurs if a model is used in an application for land-use where it was not developed. For example, SOMM model is developed for forestry/grassland and when it is used to simulate soil organic carbon in arable soils, it is forced to count grass as the only crops. That kind of structural problem causes errors because it is well documented that different plant species and management practices result in different SOC accumulation and should be considered in simulations.

Smith et al. (1997) also tested the performance of combined models and concluded that the coupling of two models does not lead to more accurate results. On the contrary, it led to inaccurate values and estimations with more errors.

Heikkinen et al. (2014) compared Yasso and RothC to measured carbon values. Soil carbon stock changes were simulated after the cropland conversion to grassland. The study area soil type was mineral soil (silt, clay and sand) and the total area was 4 hectares. The soil sampling was conducted in 1980, 1999 and 2004, and the sampling depth was 0-20, 20-40 and 40-60 cm. Soil C stock (equal to C content x bulk density) was determined to the depths 20 and 60 cm. In this Heikkinen et al. (2014) study, model values were compared to measured values using modelling efficiency (value closer to 1 indicates better performance). The concluded result was that Yasso07 and RothC both estimated the carbon stock increase relatively accurately ( $R^2$  0.60 and 0.72 respectively). Yasso07 underestimated the change in soil C stocks and RothC overestimated the changes.

## 8. Spectral methods

Spectroscopic techniques rely on the interaction of electromagnetic radiation and matter. Spectroscopic technique has many advantages over the traditional methods, as it is a non-destructive analysis method, don't require toxic or expensive chemicals, is fast, measures several parameters in a single analysis and can be used *in situ* or in a laboratory (Carcia-

Sanchez et al. 2017). Laboratory devices are accurate, but even they enable analysis of larger amount of soil samples more rapidly than dry combustion method (England and Viscarra Rossel 2018), they still require field sampling and some sample preparation before analysis, which are laborious. Proximal soil sensing is method where field-based instruments are used. It enables less time consuming and laborious estimations of soil content, and it can be used for soil organic carbon determinations in a purpose of carbon accounting (England and Viscarra Rossel 2018). Remote sensing is a method where different sensors are used from distance, for example, from airplanes or satellites. Remote sensing is used for large scale biomass estimations (aboveground carbon stocks), and although other spectral methods are typically used for below ground carbon estimations, remote sensing is not yet robust method enough for that. Remote sensing of soil organic carbon is also limited to few first centimeters of bare topsoil (Angelopoulou et al. 2019). Here the focus related to spectral methods is in soil sensing and remote sensing for above ground biomass, because of their usability and advantages.

Spectroscopic methods are utilized in several spatial scales. Spectral sensors can be used from fine to global scale. Proximal sensing ranges from micro to landscape (plot to farm) scale (England & Viscarra Rossel 2018) and remote sensing can be conducted on a regional to global scale depending on the resolution of sensors and purpose of the use (Angelopoulou et al. 2019).

## 8.1. Sensing soil organic carbon

Soil organic carbon concentration can be measured by different sensors relying on electromagnetic radiation because soil matters organic bonds and minerals absorb light in specific wavelengths. Soil information can be measured via sensors using signals that correspond to physical qualities and that information can be linked to soil properties. Soil content specific absorbance spectrum can be compared to spectral measurements of known sample via statistical model (Carcia-Sanchez et al. 2017). Reference spectrum is derived from samples, which have been analyzed by traditional laboratory methods (e.g. dry compustion) (Stenberg et al. 2010).


Soil organic carbon wavelengths are mainly in the visible-near infrared, near infrared and mid infrared region (Smith et al. 2020). Soil spectrum can be generated by directing radiation to the sample, which makes the molecular bonds to vibrate. Vibrating molecules absorb light and eventually produces a characteristic shape that can be used for analytical purposes (Stenberg et al. 2010).

## 8.2. Proximal soil sensing

According to Viscarra Rossel et al. (2011) proximal soil sensing is a method where field-based sensors are used in contact or close to (within 2 m) the soil. Proximal soil sensing doesn't include remote sensing or laboratory measurements.

There are several different proximal soil sensors and they can be classified according to the manner they measure or the source of their energy (Fig. 9). Sensors can be invasive, which means that sensor is in contact to soil during measurements or non-invasive. Invasive measurements can be done within the soil or for example excavated soils. Proximal soil sensor is active if it produces its own energy from artificial energy source and passive if they use natural radiation energy from sun or earth. Sensors can be used "on a go" or moving (mobile). Sensors can consider to be indirect or direct. Direct proximal soil sensing means that measured soil property is based on a physical process (e.g. clay mineralogy). However, when the measurement is of a proxy and inference is with a pedotransfer function (PTF = raw soil data translated into more useful information), then the proximal soil sensor is indirect. Proximal soil sensing is done in same spatial scale as conventional methods (soil sampling and dry composition) (Viscarra Rossel et al. 2011).

"Measurement of Soil Carbon Sequestration in Agricultural Systems" launched by Australian Government 2018 is the first methodology that legislates sensing for soil organic carbon estimations (England and Viscarra 2018).



**Fig. 9.** Different proximal soil sensing approaches. Measurement can be invasive or non-invasive, energy source can be natural or artificial, operation can be mobile, or stationery and inference can be indirect or direct. Closely adopted from Viscarra Rossel et al. 2011.

Soil sensors for field use are developed for cost-effective and rapid soil organic carbon determination. The benefit those sensors provide is that high sampling density can capture

more effectively field variability and hence solve the problem with selecting a correct soil sampling design (Sinfield et al. 2010).

### **8.2.1 Visible and infrared reflectance spectroscopy**

Visible (vis) portion of the electromagnetic spectrum is 400-700 nm and near infrared (NIR) region is 700-2500 nm. Visible and infrared spectroscopic techniques are sensitive to soil organic and inorganic components. Visible and near-infrared ranges can also be combined; this synergy adds value to the sensing because both wavelength ranges provide different information about soil organic carbon (England and Viscarra Rossel 2018). Mid-infrared (mid-IR) electromagnetic spectrum region is from 2500 nm 25000 nm and it contains more information about soil organic composition than vis-NIR (Viscarra Rossel et al. 2011).

Visible-NIR techniques have been used successfully in SOC concentration estimations on field conditions. Mid-IR can predict SOC concentration, most commonly, in a laboratory with measurements on dried and finely ground soil samples. This is partly because there are strong water absorptions in the mid-IR range. This effect tends to mask or deform other soil constituents' absorptions which makes it more difficult to calibrate adequately. Portable mid-IR devices are currently under development and some exist already (England and Viscarra Rossel 2018).

### **8.2.2 Laser-induced breakdown and neutron induced gamma-ray spectroscopy**

Laser-induced breakdown spectroscopy (LIBS) and neutron-induced gamma-ray spectroscopy (INS) are emerging and promising techniques (FAO 2019, Paustian et al. 2019). Laser-induced breakdown spectroscopy is a technology where optically focused short-pulsed laser is used to heat the soil sample. Heating results in the formation of high temperature plasma. After the plasma is cooled down, it can be measured with spectrometer (190-1000 nm), as plasma emits radiation that is characteristics to its fragments. The plasma forms on only a very limited area allowing only a small portion of the sample to be measured during each event. Advances in fiber optic technology also makes LIBS systems portable and mobile (Viscarra Rossel et al. 2011). LIBS measurements are rapid, in a laboratory, time per sample is less than a minute (England and Viscarra Rossel 2018).

Currently LIBS for SOC measurements are mainly conducted with benchtop laboratory devices that require sample preparation. Other constrains are sample representativeness

(tiny sample volume) and limited understanding of wet field measurements accuracy (Viscarra Rossel et al. 2011).


Inelastic neutron scattering acts in the gamma ray region of the electromagnetic spectrum. Neutron generator generates fast neutrons which penetrate the soil and interact with nuclei of the elements. In that process, gamma rays are simulated, and those rays can be detected by a scintillation detector (e.g. sodium iodide). The detector measures the spectra and from peak intensities, with specific calibrations, it is possible to determine the SOC in units of g C /m<sup>2</sup>. INS is a non-destructive method with the capability to measure to a depth of 30-50 cm, from a relatively large footprint (diameter 150 cm) and large volumes (0,3 m<sup>3</sup>) (Wielopolski et al. 2008). INS method requires some conventional soil sampling for correlation establishing, but once correlation is formulated there is no need for further soil sampling (Izaurrealde et al. 2013). In INS technique, the sampling depth is not precisely defined, but according to Izaurrealde et al. (2013) about 90% of the detected signals was from 30 cm depth and 99% effective depth was 50 cm (based on Monte Carlo calculations). This means that depth variation should not play a major role in total, since only small signal arrives from deeper layers. INS is not yet well developed, but it is known that it would suit well in field conditions.

### 8.2.3 Uncertainties and important issues

Reflected soil absorbance needs several processing steps (Fig. 9.) before information of soil organic carbon can be interpreted. Spectral analysis is based on multivariate statistical methods. Soil matrix is a complicated mixture of different parameters that have overlapping absorptions and low concentrations. These factors interfere the measurements which result to preprocessing in a purpose to for example minimize noise and enhance signals (Nawar et al. 2016). Multivariate spectroscopic modelling is a process where sensed soil properties are related to the absorbance of a set of known reference samples. This describes the relationship between spectral data and soil properties. Most commonly used method is partial least squares regression (PLSR). Accuracy of the estimations is highly dependent on the chosen calibration method (Angelopoulou et al. 2020). Model development needs several diagnostic steps, where the model fit, and performance is evaluated and improved. Model is also validated with external data set. According to "Measurement of Soil Carbon Sequestration in Agricultural Systems" methodology (2018) the data for the spectroscopic modelling and validation is divided into three components: training set, validation set and prediction set. Training set is used to develop the spectroscopic model and validation set is used to test the accuracy of model's estimates. Training and validation set's soil samples need to be analyzed with reference analytic method, which is in this case dry combustion, to determine their soil organic carbon concentration. After laboratory analysis data sets can be

used to develop the spectroscopic model. Data from large spectral libraries can be used to expand the site-specific models and to improve the accuracy, but not to develop the model.

In field conditions, soil moisture, surface roughness and vegetation cover affect the spectral signal (Angelopoulou et al. 2020). For example, vegetation would lead to overestimation of soil organic carbon (Angelopoulou et al. 2020).



**Figure 10.** Flow diagram of different procedural steps of visible-infrared spectra when estimating soil organic carbon content (Closely adopted from soil carbon methodology by Australian Government 2018).

## 8.2.4 Accuracy and cost

Stevens et al. (2006) tested field spectroscopy in agricultural land and they concluded that according to the results, the standard error of measurements allows to detect a soil organic carbon change of 7.2-9.9 Mg C/ha (upper 30 cm of the soil). According to Freibauer et al. (2004) soil organic carbon changes result of land conversion or management practices are only 0.3-1.9 Mg/ha/year, which means that the field spectroscopy would detect any changes in the soil organic carbon after minimum of 10 years. But since then, soil sensing has become an accepted method in measuring carbon sequestration in agricultural land in Australia.

England and Vicarra Rossel (2018) gathered information about accuracy and costs of different spectroscopic techniques (table 5.) for soil carbon accounting. Both LIBS and INS techniques are also expensive, sensor costs are even more than 100 000 AUD (England and Vicarra Rossel 2018).

**Table 5.** Assessment of cost and accuracy of different sensing methods for carbon determination. Modified from England and Vicarra Rossel 2018.

Method	Instrument cost (in thousands of AUD)	Measurement cost per sample (AUD)	Accuracy	
			RMSE	R <sup>2</sup>
vis-NIR, dried ground	10-100	8	0.44	0.85
vis-NIR, field condition	10-100	0.8	0.47	0.81
mid-IR, dried, finely ground	25-90	15	0.11	0.93

Cremers et al. (2001) evaluated LIBS technique's capability for soil total carbon detection. They conducted the study on agricultural soil and on woodland soil, and they concluded that LIBS instrument has a detection limit of 300 mg C/kg, a precision of 4-5% and accuracy of 3-14% (= 750 mg C/kg).

Wielopolski et al. (2011) tested INS for soil carbon pool determinations *in situ*. They concluded that INS estimations of soil total carbon amount agreed with dry combustion method values in organic soils (difference varied from 3% to 9%). In pasturelands there was no agreement between the two methods and difference varied from 16% to 168%. In organic soils, the total carbon amount varied from 3.92 to 5.36 kg C/m<sup>2</sup> and in pasturelands the variability was from -3.07 to 5.96 kg C/m<sup>2</sup>.

Izaurrealde et al. (2012) tested portable LIBS, INS and infrared spectroscopy (DRIFTS) methods against results from dry combustion. In their study, the soil carbon density (total carbon, not organic carbon) determined with dry combustion method was 4.07 kg C/m<sup>2</sup>, LIBS gave estimation value of 3.27 kg C/m<sup>2</sup>, infrared 4.32 kg C/m<sup>2</sup> and INS 2.57 kg C/m<sup>2</sup> with "universal" calibrations and 4.06 kg C/m<sup>2</sup> with "local" calibrations. Compared to the dry combustion method, the LIBS underestimated (20%) the carbon concentration, infrared overestimated it slightly (6%) and INS with local calibration was very accurate.

Different studies are hard to compare due the different measurement, modelling and prediction procedures. According to Angelopoulou et al. (2020) proximal soil sensing *in situ* is developed in recent years, but more research needs to be done.



# 9. Remote sensing

Remote-sensing procedures have been applied to collect information about above ground biomass. Vegetation structure, biomass and productivity can be estimated on a large scale by measuring the spectral reflectance of the vegetation (Vicharnakorn et al. 2014). According to Canada center for remote sensing, remote sensing refers to a science where information concerning the earth's surface is collected by sensing and recording emitted or reflected energy. Processing, analyzing and applying that information is part of that science. Remote sensing can be conducted from space (satellite systems) or air (airplane, drone). Aerial sensing is used for local-scale assessment of earth surface and satellite systems can be used for larger spatial extents (Brown 2013.). Remote sensing always needs some field measurements, the ground truthing, that is the only way to link the sensor data to biophysical phenomena. Measurements done with remote sensing instruments are physical quantities and they may or may not be useful for understanding the phenomenon of interest (Brown 2013.).

Aboveground biomass is not spatially mapped even in countries where systematic forest inventories are conducted. Mapping would add important information about carbon stocks, because forest characteristics can differ greatly from inventoried ones. Some areas are not easily accessible and remote sensing would be good opportunity for those. On a national scale, above ground carbon stocks can be overestimated or underestimated if only field approach is used (Brown 2013).

## 9.1. Theory

Remote sensing needs an energy source that illuminates or provides electromagnetic energy. This energy travels towards its target of interest and while travelling comes into contact with atmosphere. Interaction with atmosphere takes place a second time as the energy travels from the target to the sensor. After the energy makes it way to the target, they interact with each other depending on the properties of both. After the interaction, the next step is to record and collect the electromagnetic radiation. This can be done via a sensor, which is not in contact with the target, that collects the scattered or emitted energy from the target. Then the energy recorded by the sensor must be transmitted to the station where the data is processed into an image. Lastly processed images can be interpreted visually, digitally or electronically, so that information about the target, which was illuminated, can be extracted (Canada Centre for Remote Sensing 2020). All electromagnetic radiation behaves in predictable ways in accordance to the basics of the wave theory. Wavelength and

frequency are the two important characteristics of electromagnetic radiation (Canada Centre for Remote Sensing 2020).

The interaction between a sensor and the surface can be active or passive. Passive sensors measure naturally available energy. These kinds of remote sensing systems can be used when the sun is illuminating the Earth (reflected energy), and during day and night when naturally emitted energy is available (for example thermal infrared) (Canada Centre for Remote Sensing 2019). Passive sensors usually record electromagnetic waves from visible (430-720 nm) and near-infrared (750-950 nm) range of light (Zhu et al. 2018). Active sensors provide their own energy source which means that the sensor emits radiation and detects and measures the radiation reflected from the target. These kinds of systems work regardless of the time of day or season (Canada Centre for Remote Sensing 2020). Active sensors use electromagnetic waves in the range of visible light, near infrared and radio (Zhu et al. 2018).

Remote sensing for quantitative carbon estimation is challenging but offers a broader scale estimation when compared with ground-based observations. Remote sensing for carbon estimations can be separated into two major methodologies. The first is the indirect measurement of carbon and the second is a method where land cover information is integrated with observations of forest inventories. In the indirect approach gross primary production (GPP) or net primary production (NPP) is estimated with several parameters that are related to vegetation functions. Such parameters include leaf area index (LAI) and photosynthetically active radiation (PAR). Both methods can also be integrated together (Iizuka and Tateishi 2015). Airplane, unmanned aerial vehicle (drone) or satellite data can also be linked directly to soil properties or data can be used to infer vegetation changes. Vegetation change means changes in aboveground carbon, and often also changes in soil organic carbon. Remote imagery can also be used as a covariate in digital soil mapping of soil organic carbon (Chen et al. 2015).

## 9.2. Resolutions

Different remote sensing instruments have different resolutions (table 6). Spatial resolutions refer to the smallest object, pixel, that sensor is capable to detect. Spatial resolution determines how detailed a picture can be. Temporal resolution tells how often satellite passes the same spot and spectral resolution refer to sensors ability to sense different wavelengths. In spatial and temporal resolution, high resolution refers to small number of units (for example days or meters), and opposite in spectral resolution, where high resolution means that sensor can sense several wavelengths. There is usually a trade-off between

spatial and temporal resolution, which means that high spatial resolution pictures can be taken less frequently (King 2000).

**Table 6.** Satellite remote sensing resolutions (example).

	Resolution		
	Spatial	Temporal	Spectral
<b>High</b>	1-35 m	< 3 days	several hundreds
<b>Medium</b>	200-500 m	3-16 days	tens
<b>Low</b>	> 1000 m	> 16 days	5

Source: Finnish remote sensing guide (2004).

For large (global and continental) scale biomass mapping the coarse spatial resolution optical sensors, such as the MODIS, are useful because they have moderate spatial resolution, and good image coverage and frequency in data acquisition (good trade-off between those). Smaller (local to regional) scale biomass mapping needs finer spatial resolution instruments to achieve data with more details (Lu 2006).

According to Lu (2006) vegetation estimations with coarse spatial resolution data over larger areas have been limited by the errors caused by mixed pixels, and the major difference between the pixel size of the satellite and the ground reference data. Mixed pixels case a situation where the coarse resolution pixels receive response from several objects (such as trees), and from that data, biomass cannot be directly estimated. Because coarse imaging satellites have useful characteristics (e.g. good image coverage), finer spatial resolution satellite data has been used to combine ground reference data to this coarser spatial resolution data. This is usually done by regression techniques (Muukkonen and Heiskanen 2006). For example, Häme et al. (1997) derived regression models from ground reference data and Landsat satellite data which they utilised successfully in the medium coarse spatial satellite data. Finer spatial satellite data models can be used as an intermediate step between ground measurements and coarse resolution data.

## 9.3. Optical sensors

Optical imaging sensors operate in the visible and reflective infrared ranges. Aerial photography was the earliest version of remotely sensed data and it has been used to local-scale assessments of the Earth's surface properties (Brown 2013). Larger spatial scale remote sensing with satellite platforms and optical systems have been used for several decades (Brown 2013). Optical instruments on space platforms typically include panchromatic systems, multispectral systems and hyperspectral systems. Panchromatic system is sensitive to broad wavelengths range and the image it provides is black and white or grey scale. A multispectral sensor detects multiple channels with few spectral bands and each of those channels are sensitive to certain wavelength. Resulting image includes the brightness and spectral colour information. A hyperspectral sensor uses information from 10 to 100 of spectral bands and one image consists of a set of images. These optical systems differ from each other, and when the light is split into multiple spectrums it lowers the resolution of the image. That's why panchromatic system has a higher resolution than a multispectral or hyperspectral image (Zhu et al. 2018).

Landsat is one of the optical satellite systems which can be utilized for carbon mapping. Landsat land observation was launched in July 1972 and it has been orbiting earth since then (Brown 2013). Landsat is high spatial resolution satellite (30 m resolution) and its collected data is publicly open. There are also commercial high spatial resolution satellites, like QuickBird (measuring visible to infrared region). Landsat and QuickBird are passive optical sensor systems. Landsat collects data at specific multiple spectral wavelengths (optical multispectral remote sensing), measuring visible spectrum. Optical instruments can also collect data across the entire spectrum of reflected solar energy (optical hyperspectral remote sensing). For example, NASA's operating airborne Visible/Infrared Imaging Spectrometer (AVIRIS) sensor is hyperspectral sensing system (Brown 2013). Satellite optical imaging system's spatial resolution varies from under 1m to 2 m (high spatial resolution) (Zhu et al. 2018). Variation depends from the sensor (e.g. QuickBird under 2 m, MODIS 250m) (Zolkos et al. 2012).

Optical sensors and their spectral measurements have been used to model and monitor primary production of above ground vegetation (Brown 2013). Soil organic carbon mapping from space via satellite imagery is usually done with multispectral images of medium or high spatial spectral resolution or hyperspectral Hyperion image. To estimate soil organic carbon content from the reflectance values in the available image bands it usually relies on multiple linear regression equation. There are several ways to estimate reflectance values from the satellite data. Usually the magnitude of the error contributing reflectance prediction is not specified (no validation with additional reflectance measurements). This reflectance

predictor error affects negatively to the accuracy of the soil organic carbon content estimations (Vaudour et al. 2013).

Optical remote sensing systems are sensitive to optical properties and moisture (Zolkos et al. 2012). Optical remote sensing does not directly assess the aboveground biomass (Vicharnakorn et al 2014), but it gives two-dimensional information, which can be linked to biophysical characteristics of the vegetation. This linkage is indirect and optical systems can be used to identify horizontal variability and, for example, to assess canopy conditions. The electromagnetic energy that optical sensors utilize is emitted or absorbed on the upper layers of vegetation, so it only gives limited information (does not penetrate through vegetation to the Earth's surface) (Zolkos et al. 2012).

Optical remote sensing means passive sensing of visible and near-infrared reflectance from the earth (for example GoogleEarth). This remote sensing system has been widely used to link direct aboveground measurements to satellite observations due to fact that different canopy structure gives different reflectance. This method is not consistent over large areas because rapidly varying surface conditions cause artefacts to the derived maps, as the satellite observations cannot keep up (cloud free time, repeat time). Frequent repeat measurement sensors, like the Moderate Resolution Imaging Sensor (MODIS), have helped to overcome this problem (Goetz et al. 2009).

## 9.4. Synthetic aperture radar

Synthetic aperture radar (SAR) is an active system that produces images of land surface based on principles of radio detection and ranging (RADAR often synonym for SAR) (Goetz et al. 2009). SAR uses microwaves, an electromagnetic spectrum range of 1 mm-1 m (Zhu et al. 2018). SAR can penetrate through haze, smoke and clouds, and it can operate during day and night. SAR transmits microwave energy which can penetrate forest canopies. SAR sensors are sensitive to different aboveground biomass components according to the wavelength of the sensor. Shorter wavelengths are more sensitive to leaves and small branches, and longer wavelengths are more sensitive to stems and large branches (Goetz et al. 2009). SAR only shows the geometry and surface roughness of the target and it does not produce data where you can identify for example the vegetation type (like infrared, and they are used to complement each other's) (Zhu et al. 2018). Several radar satellites are currently operating, for example the European ENVISAT/ASAR, the Japanese ALOS/PALSAR and German TerraSAR-X (Goetz et al. 2009).

One of SAR system's disadvantages is that its estimations of AGB are limited as the SAR instruments lose their sensitivity with increasing biomass. This phenomenon is known as

“saturation” and it occurs in relatively low, but undetermined, biomass densities (optical saturation point is around 100-150 Mg/ha and for SAR a bit higher) (Zolkos et al. 2012). Low saturation point causes uncertainties in AGB mapping because estimations of vegetation density in high density forest is not accurate.

Current SAR systems can produce an image with a half meter of accuracy (Zhu et al. 2018). It measures forest structure with high spatial resolution (20-100 m), can operate regardless of the time of the day and it can penetrate through clouds and through vegetation (Zolkos et al. 2012).


## 9.5. Lidar

Lidar uses a pulse of energy from a laser operating at optical wavelengths to actively sense vegetation. Lidar systems that are typically used for vegetation mapping usually operate in wavelengths between 900 and 1064 nm. They record the time the pulse is travelling, and that time-return interval can be used to calculate distance between the sensor and the object (Zolkos et al. 2012). Lidar’s laser beam width varies (small to large footprint) and a small footprint beam typically illuminates a surface area with a diameter 50 cm or less. This kind of accuracy provides information that is increasingly utilized in forestry applications but in a small area. Medium to large footprint lidars illumination surface area can vary from larger than 5 m to approximately 65 m (depending the platform) (Zolkos et al. 2012). A lidar can measure the three-dimensional vertical structure of vegetation in great detail (Vierling et al. 2008). This information can be applied to above ground biomass via correlative models which has been derived from associated field measurements (Zolkos et al. 2012).

There are some lidar’s operating form satellite platforms. The geoscience Laser Altimeter System (GLAS) spaceborne lidar system which be used to estimate forest aboveground biomass on a large scale (Xiaofang et al. 2019). Airborne lidar is scalable and cost effective (Asner et al. 2013). Lidar and biomass have reported to show strong relationship beyond biomass levels of 1000 Mg/ha, which is far more than SAR and Optical sensors are capable. Lidar can estimate the vegetation structure direct (e.g. canopy hight distribution) and it has been shown in several studies that it provides more accurate results in AGB estimations than optical and SAR data. Lidar can’t penetrate clouds (Zolkos et al. 2012).

## 9.6. Important issues influencing biomass estimations for carbon mapping

Remote sensing is an indirect way to estimate aboveground biomass and there are several critical steps to consider when building a proper biomass estimation procedure (Fig. 11). The selection of the sensor and platform, sample size of the reference data, variables, algorithms and cross validation of data is important, and each step include uncertainties (Xiaofang et al. 2019). Many studies have conducted aboveground estimations, but comparing those studies is difficult due to diversity of data sources, modelling methods and calculation standards.



**Fig. 11.** Workflow of satellite image processing. Combination of Sentinel and Landsat data. Closely adopted from Xiaofang et al. 2019.

Assessing carbon stocks with remotely sensed data uncertainties are high. Each workflow step includes possible error sources and ecological subjects are hard to monitor with high accuracy. This means that vegetation structural variations, species composition, heterogeneity of landscapes, soil properties, climatic and topographic variables and disproportionate data availability all create high uncertainties that affect biomass division and change tendency (Issa et al. 2020).



## 9.6.1 Remotely sensed data

Remotely sensed data introduces limitations and sources of uncertainties. Different sensing systems provide different information and the selection of the right sensor data is essential. For example, each sensor type provides different resolutions and different information according to the polarization and angularity (Lu 2006). Sensor selection according to a specific purpose and study area is the key, and since each sensor type has their own characteristics, they can be integrated to achieve data where one sensor's limitations are exceeded. Multisensor synergy can produce estimations with accuracy levels similar to those of the lidar alone, or even better accuracy, but study results have varied (Zolkos et al. 2012).

Remotely sensed data needs also several corrections due to the radiometric characteristics and interaction with the atmosphere (Lu 2006). When electromagnetic energy passes through the atmosphere it causes changes to direction, intensity and spectrum of the radiation. These atmospheric effects need to be corrected, and typically it is done via mathematical models. Without proper atmosphere corrections, satellite pictures will show major distortions (Muukkonen and Heiskanen 2006). Atmospheric correction with mathematic models is necessary, but they also include inaccuracies (Muukkonen and Heiskanen 2006). Topographic factors also influence the reflectance of the vegetation and remotely sensed data from mountainous regions needs removal of topographic effects (Lu 2006). It goes without saying that remotely sensed data handling requires a thorough knowledge of data processing and an understanding of the phenomena under interest.

## 9.6.2 Reference data quality and variables

Remote sensing data is conventionally compared with forest *in situ* measurements (non-destructive estimations). Forest field measurements include tree height, stem diameter and density. These *in situ* measurements can be converted to forest biomass estimations by allometric models and they are usually conducted on a local scale (Zolkos et al. 2012). Forest field measurements are essential for biomass estimations and field measurements can also be called reference data. The field measured data can be used for different purposes e.g. model development, validation, calibration, comparing different models and to conduct uncertainty analysis. High quality data source is thus essential for developing an AGB estimation model. The quality of the field data may vary greatly because data is essentially collected for another purposes, tree species composition might be very complex and wood density may differ (Lu 2006). Quality of field data affects the accuracy, and calibration or validation of the calculated AGB is needed. Asner et al. (2013) argued that to reduce uncertainties in lidar and satellite measurements, the necessary step is to measure plot-level

biomass instead of estimating it from conventional inventories with allometric equations. More accurate plot level direct measurements offer better data for lidar calibrations.

Geometric accuracy of the field data sample plots and remotely sensed data is also important factor affecting the accuracy of the biomass estimations (Lu 2006). Without proper geometric accuracy, the relationship between AGB and remotely sensed data can be erroneous.

One factor affecting biomass estimations is selection of suitable remote sensing variables. Remote sensing variables, such as spectral signature, vegetation indices and image textures, may be suitable parameters to describe AGB. Suitable variables should correlate significantly with AGB and weakly with each other's. Weak correlation with AGB reduces the AGB estimation performance. Potential parameters can be identified with several statistical methods. For example, stepwise regression analysis and correlation analysis can be used. Both are simple analysis which are based to the relationship between AGB and tested variables (Issa et al. 2020).

### 9.6.3 Modelling and uncertainty analysis

Different modelling algorithms can be used to describe what remotely sensed data tells about phenomena of interest, which is in this case aboveground biomass (carbon pool). Algorithms link reference data to the variables divided from remotely sensed images. Empirical algorithms include parametric and nonparametric algorithms, and both are widely used in aboveground biomass estimations. Parametric algorithms (e.g. simple or multiple linear regression model) and nonparametric algorithms (e.g. K-nearest neighbors, random forest) behave a bit differently. Parametric algorithms assume straight forward linkages between variables and biomass and estimation are based on models which predict the relationship. But in real life, factors affecting biomass are complex and numerous, which means that it is difficult to predict the relationship with simple regression models. An alternative approach is to use nonparametric algorithms where the model structure is predicted in a data-driven manner, instead of explicitly predefining the model structure (Xiaofang et al. 2019).

Several factors determine which model/algorithm/prediction method is selected. Factors, such as, sensors resolution, the availability of biomass sample data and reference data, the scale of the area, the availability of related software and human resources all affect the selection of prediction model. Different models perform differently according to factors mentioned above. One way to evaluate model performance is uncertainty analysis. Uncertainty analysis is used to assess the accuracy of biomass estimates. The root mean

square error (RMSE) and the coefficient of determination ( $R^2$ ) are commonly used measures (Lu 2006). High  $R^2$  value and low RMSE value often indicates that developed model fits well with the sample plot data. Traditionally, in remote sensing data analysis, the reference data is split into two parts, where one part is used to develop the model, and other is used for evaluation of model performance (cross-validation). Uncertainty analysis is one way to compare performance of different approaches but there is also variation between analyzed results. For example, one of the common types of cross-validation method is k-fold cross-validation (Xiaofang et al. 2019), but the value of k differs in different studies and that might affect the model diagnostics.

Xiaofang et al. (2019) studied the performance of different prediction methods by combining Lidar data (GLAS), optical data (MODIS) and field measurements. They compared six prediction methods (Gaussian processes, stepwise linear regression, nonlinear regression, partial least squares regression, random forest and support vector machines) and effect of prediction method, sample size of field measurements and cross-validation settings. Authors concluded that prediction method had the most considerable effect on the quality of the estimations. In most cases the random forest -model produced more accurate predictions than others. The sample size obviously affected the prediction model performance and for example the random forest algorithm combined to large number of field measurements (n=801) gave the most precise results ( $R^2=0.73$  and  $RMSE=23.58$  Mg/ha).

Several approaches have been developed, yet there is no universal model, or at least agreement which model would perform best in certain situations. One universal model might be too difficult to develop (Lu 2006), but it is necessary to identify (or develop) models suitable for different environments. More advanced models for AGB estimations, which utilize multi-source data, are also needed.

Asner et al. (2013) demonstrated the uncertainty of aboveground carbon density estimations by combination of Lidar sensors and satellite (Landsat) data. Accuracy was compared to field measurements. Study was conducted in Panama, and study site was the whole country. Vegetation types ranked from dense tropical forest to grasslands. The result demonstrated that the lidar based carbon mapping has an uncertainty of about 10% at 1 ha resolution.

Fassnacht et al. (2014) compared how sample size, sensor type and prediction method affect the accuracy of the AGB estimations. They conducted the study in two locations, in Europe and South- America, and they compared three different sensor scenarios and they performance with different sample sizes and different prediction methods. Authors concluded that selection of sensor type had the highest impact on accuracy. In their study, the best performing sensor scenario was a combination of airborne lidar data and spaceborne optical data (compared to airborne optical and airborne lidar). Best performing prediction method was again random forest. The overall best performance was multisensory

synergy with random forest algorithm and biggest sample size ( $R^2$  0.71 and RMSE 37, case mean was  $R^2$  0.42 and RMSE 52). The main discovery in their study was that a good prediction method might be more important than increasing the number of the field data.

## 9.7. Accuracy

Accuracy requirements for remote sensing approaches are not yet explicitly stated, but studies conducted by Hall et al. (2011) and Houghton et al. (2009) asserted that satellite remote sensing should give errors in biomass estimation within 20 Mg/ha or 20% of field estimates. Errors should not exceed 50 Mg/ha for a global biomass map at 1 ha resolution (Zolkos et al. 2012). Asner et al. (2013) concluded that fraction cover of photosynthetic and non-photosynthetic vegetation imagery from Landsat, combined to topography and climate data, is a suitable way to map national-scale aboveground vegetation density on a per hectare basis. This method gives low uncertainties to estimations. In Panama study, Asner et al. (2013) calculated that carbon density uncertainty was on average 20.5 Mg C/ha at the national level.

## 9.8. Cost

Economic condition is probably the most important factor affecting the implementation of satellite remote sensing for carbon mapping. Economic condition influences the extent of the field work, purchase of different sources of image data and the human resources that all affect the accuracy of the AGB mapping. Higher accuracy often means higher costs. Asner et al. (2013) estimated that their costs for airborne lidar inventory was 1 USD dollar (0.85 €) per hectare in an area of 600 000 hectares. Costs of airborne lidar approach decrease when area increases and Anser et al. (2013) estimated that for example cost for twice as large area, the cost per hectare (airborne lidar acquisition and analysis) would be about 0.15 USD dollars (0.13 €).

Böttcher et al. (2009) collected costs for monitoring Reduced Emissions from Deforestation and Degradation (REDD) projects in different areas and different project scales (table 6.). They concluded that cost can vary from 0.42 to 463 €/km<sup>2</sup>. These project areas were smaller (40-28000km<sup>2</sup>) than area in Asner et al. (2013) study, but results differ from each other largely.

**Table 6.** Biomass monitoring costs with different remote sensing techniques (acquisition and analysis costs) example.

Satellite, sensor	US\$/km <sup>2</sup> (€/km <sup>2</sup> )
Optical, medium	0.50-1.21 (0.42-1.02)
Optical, high	7.50-35.40 (6.32-29.83)
Optical, very high	116-272 (97.74-229.18)
Radar, satellite SAR	7.04-10.54 (5.93-8.88)
Radar, airborne SAR	>345 (290.68)
Lidar, airborne	100-550 (84.26-463.41)

## 10. Gas flux

The full carbon budget on the ecosystem level can be achieved with carbon flux measurements (Smith et al. 2020). Micrometeorological techniques measure the gas exchange between ecosystem and atmosphere (Rinne et al. 2016). Full carbon budget quantification needs information about carbon uptake through photosynthesis, carbon losses through respiration and other C inputs and outputs. Gross primary production (GPP) describes the carbon uptake through photosynthesis and when soil, plant and litter respiration are subtracted from the GPP, the result gives the net ecosystem exchange (NEE), or net ecosystem production (NEP). Net primary production (NPP) describes the amount of photosynthates that are available for plant growth and other functions. NPP links GPP and respiration (equation 10) (Smith et al. 2010).

$$GPP = NPP + R_a \quad (10)$$

where

GPP = gross primary production

NPP = net primary production

R<sub>a</sub> = autotrophic respiration.


NEE and NEP can be estimated with cuvettes (not included) or with the eddy covariance method.

## 10.1. Eddy covariance

The eddy covariance (EC) is a method for measuring heat, mass and momentum exchange between the surface and the overlying atmosphere. Surface should be flat and horizontally homogenous, and under these conditions the net transport is one-dimensional. The vertical flux density can be calculated by a covariance between turbulent fluctuations of the vertical wind and the quantity of interest (Aubinet et al. 2012). This means that the exchange rate of CO<sub>2</sub> between atmosphere and ecosystem can be determined with measuring the covariance between the vertical wind velocity fluctuations and the CO<sub>2</sub> mixing ratio (Byrne et al. 2007). An eddy is a current of flowing material that moves in a whirlpool or circular motion against the main current. Eddies are formed from wind, roughness of terrestrial surface and convective heat flows on the boundary layer. A boundary layer is a relatively thin layer of the atmosphere and the thickness varies diurnally and geographically (Posudin 2014).

The eddy covariance system includes three sensors that are used to measure water vapor density, air temperature and wind speed. EC measurements are typically made on the surface boundary layer. Fluxes are approximately constant in height on the surface layer; hence measurements taken in this layer are representative of the fluxes from the underlying surface. On the boundary layer the atmospheric turbulence is the main transport mechanism. To determine CO<sub>2</sub> fluxes the tower needs also analyzer for measuring turbulent fluctuations in CO<sub>2</sub>. Commonly used analyzer is an infrared gas analyzer, with open- or closed-path configuration. Open-path system needs more maintenance above cultivated soil, because for example dust from tilling can block the equipment. Both systems include a broadband infrared light source, band-pass filter and a detector. When infrared light is emitted by CO<sub>2</sub> molecule the detector observes the reduced light intensity. This reduction presents the nonlinear function of the molar concentration of CO<sub>2</sub>. To be able to capture eddies the air movement should be constant. Eddy covariance sampling frequency is high, usually 10-20 Hz. This is because only high sampling frequency can cover turbulent fluxes (Aubinet et al. 2012). Eddy covariance measurement tower can cover large scales from hundreds of meters to several kilometers or it can be used in field scale. This sampling area is called flux footprint, and size of it depends from the objective of measurements (Byrne et al. 2007).





**Fig. 12.** Eddy covariance tower and power source in a grass field. Photo: Anniina Lampinen

### 10.1.1 Important things to consider

Application of this micrometeorological technique requires vast knowledge of technical issues and the studied phenomena. There are several options for eddy covariance tower design, and some instrument options and configurations. Towers can also be placed on different sites. Installation and operational costs, and precision and accuracy are the main points taken into consideration when designing the optimal solution and selecting among the options. Site design is the first step to ensure proper accuracy and precision. There is unlikely going to be a perfect design and some compromises between science requirements, costs and practicality inevitably occur. The measurement tower should be placed in an area where the systematic biases are minimized, and ecological integrity is maintained. Ecosystems are structurally and functionally diverse, but all the environmental extremes can be found. Towers need to capture complex ecological drivers and processes and withstand extreme conditions like high temperature, ice and snow loading. After site is selected, site operation including calibration and validation are essential steps to make sure



that overall performance is as good as possible, and associated uncertainties are minimized (Aubinet et al. 2012).

With ecosystem gas flux measurements, the effect of diurnal, seasonal and annual variation should be considered. According to Flechard et al. (2020) if eddy covariance flux measurements are used to quantify the amount of carbon that is accumulated in forests the study period should be one or several decades. This is because only long measurement period would ensure statistically significant results, because the annual stock changes are relatively small compared to the large carbon forest stock, especially if forest soil is included.

### 10.1.2 Uncertainties and accuracy

Eddy covariance methods includes several possible sources of uncertainties. According to Baldocchi (2003) when operating the most accurate results can be measured when the tower locates in flat terrain on top of homogenous vegetation which extends upwind for an extended distance surrounded by steady atmospheric conditions. But in nature conditions are not always ideal and when the method is applied over complex and natural landscape where wind, temperature, humidity and CO<sub>2</sub> vary with time, the CO<sub>2</sub> quantification needs corrections. Long measurements period reduces the random sampling errors, but long measurement time means also gaps in the data. Gaps in a long-term data can occur for example due to sensor malfunction or if the wind is coming from undesirable wind sector. Gaps in the data can be filled e.g. with empirically derived algorithms. This approach needs continuous tuning because biological factors (like leaf area, soil moisture) are changing seasonally. Nighttime fluxes need also corrections. CO<sub>2</sub> emitted by nighttime may not reach the tower height due the fact that during nighttime thermal stratification is stable. If nighttime CO<sub>2</sub> is not measured, then the system underestimates the ecosystem respiration.

Goulden et al. (1996) concluded that sampling error in forest study area was  $\pm 30$  g C/m<sup>2</sup>/year and the net annual CO<sub>2</sub> uptake is 200 g C/m<sup>2</sup>/year. According to Baldocchi (2003) the error bound from nearly ideal sites for annual net exchange of CO<sub>2</sub> is less than  $\pm 50$  g C/m<sup>2</sup>/year (concluded from several studies). Carbon sequestration estimations from flux measurements have been relatively uncertain.

# 11. Other methodologies

This chapter provides short overview of quantification methods used in current offset projects by two companies that share the majority of all offsets in voluntary markets. Purpose is to introduce quantification methods these companies use and how the uncertainties are considered, to give perspective of current state.

There are several companies, acting in a voluntary market, that provide carbon offsets via different projects and activities. This area is not currently regulated, and it lacks standards and specific project boundaries (e.g. what areas can be used for afforestation). This have led to the situation where individuals or companies interested about compensating their CO<sub>2</sub> emissions do not know how all the difficult issues associated to carbon sequestration (see section 3.7.1) are dealt in practise. Some companies or non-profitable organizations provide methodologies which aim to ensure the actual positive and sustainable environmental effects are achieved. After the compensation project have been carried as describe in the provided methodology, the company can certificate those offsets. According to Financial news (2019) Verra and Gold Standard are companies certifying 80% of all offsets. Both companies have been launched in beginning of 2000s and aim to sustainable climate actions. Companies have developed several methodologies for different natural based actions and most deposited and released credits are from forestry practices (afforestation) which are implemented in developing countries. Each methodology includes descriptions of how different part of projects/activities are dealt.

## 11.1. Verra

Verra is a global company that develops and manages standards for sustainable development and climate actions. Their standards and frameworks help to channel finances to projects and activities which have high positive climate impact. Verra was founded in 2005 by specialists to fill the need for more assurance quality practises in voluntary carbon markets. Nowadays Verra manages several programs and initiatives, like VCS Program, where they turn greenhouse gas emission reductions and removals from certified projects into tradable carbon credits. This program includes different technologies and natural based solutions, including forest and wetland restoration and conversation, and agricultural land management (Verra 2020).

They have developed Agricultural Land Management (ALM) methodology which was launched in October 2020. This methodology provides procedures to estimate the GHG (CO<sub>2</sub>,

CH<sub>4</sub>, and N<sub>2</sub>O) emission reductions and removals which are achieved with improved agricultural land management practises. Focus is on practises that help to increase the soil organic carbon storages and it can be used in a regenerative agriculture. ALM methodology provides three possible approaches:

1. Measure and Model:

Initial SOC stock size is measured and an acceptable model is then used to estimate the change. Acceptable model needs edaphic characteristics and actual implemented agricultural practices, initial SOC stock and climatic conditions from sample fields.

2. Measure and Re-measure:

This approach utilizes direct measurements in two time points, and it can be used if there are no relevant models to apply. Some regions, crops or practices may lack accurate validated and parametrized models. In October 2020 it is mentioned that this quantification approach 2 cannot be used because there is no benchmark for performance.

3. Calculation:

CO<sub>2</sub> flux is calculated from IPCC Guideline for National Greenhouse Gas Inventories (2019) equations.

ALM methodology concludes that in their projects the key sources of uncertainties are sample error, measurement error of model inputs and error in model prediction. They include uncertainty reduction value to results and it can be calculated with equation 11.

$$UNC_t = MIN(100 \%, MAX\left(0, \frac{T \sqrt{\sum s_{\Delta,t}^2}}{\Delta CO_{2t} + \Delta CH_4t + \Delta N_2O_t} - 15 \%\right)) \quad (11)$$

where

UNC<sub>t</sub> = Uncertainty deduction in year t (expressed as the extent to which the half width of the 95% confidence interval, as a percentage of the mean, exceeds the threshold of 15%); unitless number between 0 and 1

∑ = Sum over pools and gases CO<sub>2</sub>\_soil, CTREE, CSHRUB, 24 CH<sub>4</sub>\_SOC, CH<sub>4</sub>\_ent, CH<sub>4</sub>\_md, and N<sub>2</sub>O\_soil, where Approaches 1 or 2 were employed.

s<sup>2</sup><sub>Δ,t</sub> = Variance of the estimate of Δ · t. (Δ · t = mean emission reductions from gas and pool · at time t) (see); (t CO<sub>2</sub>e/unit area)<sup>2</sup>

ΔCO<sub>2t</sub> = Areal average carbon dioxide emission reductions in year t; t CO<sub>2</sub>e/unit area

ΔCH<sub>4t</sub> = Areal average methane emission reductions in year t; t CO<sub>2</sub>e/unit area

ΔN<sub>2</sub>O<sub>t</sub> = Areal average nitrous oxide emission reductions in year t; t CO<sub>2</sub>e/unit area

$T$  = Critical value of a student's t-distribution for significance level  $\alpha = 0.05$  (i.e., a  $1 - \alpha = 95\%$  confidence interval) and the degrees of freedom  $df$  appropriate for the design used (e.g.,  $df = n - 1$  for a simple random sample of  $n$  sample units)

15% = Threshold beyond which there is an uncertainty deduction

• = Gas or pool

(Verra Agricultural Land Management methodology 2020)

## 11.2. Gold Standard

Gold standard for the Global Goals is standard that provides credible projects that have positive effect to climate and sustainable development. Gold Standard was established in 2003 by WWF and other non-governmental organizations. Paris Climate Agreement and the Sustainable Development Goals were the main drivers to develop a best practice standard for climate and development in a most sustainable way. Their purpose is to maximize positive climate impact and at the same time providing income possibilities for people. Their mission is: *"To catalyzes more ambitious climate action to achieve the Global Goals through robust standards and verified impacts"* (Gold Standard 2020), which means that they really emphasize that under their certification, results are trustworthy

Gold Standards for the Global Goals launched Soil Organic Carbon Framework Methodology in January 2020. This methodology introduces requirements for soil organic carbon stock and greenhouse gas emission quantification. The methodology is exerted to agricultural lands and SOC changes through better agricultural practices can be quantify with this. Gold Standard SOC methodology takes into account that data or measurements needed are not available in all projects or activities, and that's why they provide three possible approaches:

1. Direct measurements (soil samples) and directly documented baseline and size of the SOC stock.
2. Data from peer-reviewed scientific publications (modelling).
3. Default factor from the IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2019) and Tier 1 and 2 approach.

With these possibilities they conclude that this methodology can be applied in a broad range of activities and with different recourses. This methodology also takes into account that science around SOC impact and activities are constantly evolving and it is not limited. Gold Standard SOC methodology sets the accuracy requirement to be precision of 20% of the mean (90% confidence level) of total SOC change calculations. The methodology includes equations how the uncertainty of SOC change models can be calculated and if the

uncertainty is bigger than 20% the project owner should include uncertainty reduction. This reduction value can be calculated with equation 1.

$$UD = UNC - 20\% \quad (12)$$

where

UD = uncertainty reduction (%)

UNC = model output uncertainty (>20%) (%).

(Gold Standard, Soil organic carbon framework methodology 2020)

## 12. Calculation examples

### 12.1. Forest stands

Increasing carbon capture in Forests: Case South-Savo

From the silvicultural methods especially fertilization and forest breeding are effective in increasing the forest production and carbon accumulation to forests (additionality) (Saarsalmi et. al. 2012).

Here we analyzed the effects of silviculture on carbon capturing by the means of scenario analysis.

The aim of the calculations was to produce an estimate of carbon capture potential of an average forest farm (32 ha) located in South Savo.

In the scenario analysis done on county level the carbon capture potential in different types of forests and different silvicultural practices was made and transformed to represent a typical forest farm.

The forecasts were done by applying the Motti calculation method developed in Luke and based on the NFII measured stand data (Hynynen et. al. 2014, Salminen et. al. 2005).

The scenarios used were

**BAU** Silviculture according to present standards

- Present level of silvicultural operations and harvested amounts
- Operations are continuing at the present level

**CARBON** Silviculture promoting carbon capture

- The harvested amounts are kept as in BAU.
- Carbon capture is increased by fertilization and expanding the rotation periods

The county level results were transformed to represent an average forest farm. The scenarios represented a 50 year period. The carbon storage consisted of living and dead trees and soil carbon. The NFI 11 (2009-2013) data consisted of 4084 experimental plots representing an area of 1,1 Mha.


The Motti simulations were done using several different chains of operations.

From the optimal solution of each scenario we got the stand characteristics (volume, biomass etc.), the amount of harvested timber and the areas of different forestry operations.

The biomass of trees was estimated with biomass equations, and biomass is transferred to carbon with 1:2 ratio (biomass carbon content 50%), carbon was changed to carbon dioxide equivalents (CO<sub>2</sub>-ekv) in the ratio of atomic weights (44/12).

The change in the soil carbon was estimated as in GHG reporting. The initial soil carbon storage is expected to be on an average level of South Finland. Estimates were made by Yasso07 –soil model.

## Simulations



**Fig. 13.** Examples of volume growth response to fertilization in two forest stands Myrtillus and Vaccinium type forests. The effect of fertilization (blue bars) can be seen in both stands.

## Calculations at two experimental farms

In co-operation with Forest Center we have calculated scenarios of fertilized/nonfertilized treatments on two selected farms, one privately own farm in South-Finland and one co-operative in Northern Finland. In addition to that, preliminary calculations for the larger set of farms we done. The analysis will be further developed in making different types of selection criteria of the stands to be fertilized and modifying the fertilization response functions according to recent data.

## 12.2. Agriculture

Carbon Farming is a new way of farming to sequester carbon in the agricultural soil. Carbon that otherwise ends up as CO<sub>2</sub> in our atmosphere, causing climate change. There are many ways to do this: from small adjustments on farm level - like applying fertilizers rich in carbon, reduced or no-tillage, or planting cover crops - to changes in the entire farming system - like enriched crop rotation or agroforestry.

In this task the main aim is to estimate the soil organic carbon (SOC) stock change for mineral agricultural soils in Finland. We applied Yasso07 soil model in this study, which describes the decomposition of organic matter based on information on climate and C input quality (Tuomi et al 2011, Tuomi et al. 2008). Yasso07 model is widely applied to assess C balances of forest and agricultural soils. It is used in the greenhouse gas inventory of Finland to assess changes in soil C stocks (Statistics Finland 2019).



Cover crops are crops planted after the harvest of the main crop, to prevent the land to be fallow. They fix additional carbon from the atmosphere by photosynthesis and offer additional biomass to the soil. They protect soils against erosion, can break infections with soil borne diseases, increase infiltration of water, fix nutrients and might increase agrobiodiversity and the overall resilience of agricultural systems.

When growing a wider diversity of crops and perennial forage crops, a more diverse agroecosystem is created. With increased diversity of soil life, roots and improved soil structure as a consequence. Such soils have a greater ability to store carbon. Introducing less intensive crops, such as cereals and grass and clover species in the crop rotation, increases the carbon content in the soil through the extensive rooting system.

We have made example calculations on SOC stock changes on crop husbandry, cattle, pig and sheep farms. We have assessed changes in hectare-based SOC stocks as a consequence/result of changing farming practices at farm level. In this approach, we compare the impact of “improved farming practice” on SOC stock to the present farming practices.

To estimate changes in the SOC stock, Yasso07 soil model uses data on climate and carbon inputs, in this case data on plant and manure inputs. To estimate the plant C input, we utilized data from farms regarding crop rotation, crop yields and cultivation area and complement data from literature. Crop residues are assumed to be left in the soil which is the normal way in Finland. C input from the residues of cultivated crops are calculated and divided to C inputs from aboveground biomass, root biomass and rhizodeposition.

To estimate C input from manure, we utilize data from farms regarding field area per animal and number of animals.

Model initialization was carried out by assuming the soil to be in a steady state with average C inputs from plant litter and manure currently used on farms. Average climate of 1990-2004 was used in the initialization. One of the decisions to be made is how long past do we go, when selecting the previous climate before the starting point of the calculations.

The effect of the climate data on the simulated SOC stock changes was studied by comparing simulations performed with different climate input data. We had a case with no climate impact in which the model initialisation and simulation were done with same climate data (period 1990-2004). The comparison was done using same climate data in the initialisation and different climate data (2005-2018) in the simulation phase

We selected a case in which farming practices continue similarly as currently as a baseline.

In our examples a 100 yr. period was selected in the simulation. During this period climate is expected to get warmer. When analysing the results, it has to be decided how this temperature change is taken into account in model calculation, because it has effects both on growth and decomposition.

When the model calculation for carbon market use is done on annual basis, the actual climate can be applied.

When selecting the model to be used it has also to be decided, do we include other climate parameters than temperature, i.e. precipitation, drought periods and the variation of them. These selections have an effect in addition to growth and decomposition also on the emergence of e.g. pests.

At present the model applied has an effect of climate only on the decomposition, not yield

Selection of the species to be grown and species rotation to be used has to be made and areal differences in the interval of species rotation applied taken into account.

More data is needed of the yield (and decomposition) of catch crops (below and above ground) and the amount of growth reduction to main crop, because the additionality of catch crops as such is obvious.


If the calculations are done in short time intervals, we have to characterize in addition to the short time variation in climatic variables also the variation in root litter and root exudates and define what is the rotation period of the roots.

The model we applied in the example calculations was Yasso, but also other soil models need to be tested in future.


The possibilities to parametrize the models are numerous, and by different parametrization we can get a wide range of outputs.

The actual processes (e.g. decomposition, species involved in that but also plant growth responses) are still poorly known and understood.

As a first impression it has shown out to be reasonably challenging to get the farmers to give a permission to use their farm as an experimental farm for calculations. While contacting the farmers to join our project, we have also discussed how this type of instruments could influence on their willingness to take carbon sequestration as one of their products.



**Fig. 14.** The annual carbon inputs (kg C ha<sup>-1</sup>) of different cultivated plant species and manure in a pig farming. Practice A represents inputs from current farming practices and practice B from improved farming practices including cover crops in the crop rotation. Aboveground C inputs consist of crop residues left on the soil after harvesting (stubble). Belowground C inputs consist of roots and root excretion.



**Fig 15.** Simulated soil organic carbon changes. The red line represents a situation in which the current farming practices continue during the simulation period and climate changes. In this case, as the C inputs do not change, the soil C stock declines due to changes in the

climate which accelerates decomposition. The Yellow line represent a situation in which the current farming practices continue during the simulation period and climate does not change. The blue line represents a situation in which the farming practices change from current to the improved farming in the beginning of the simulation period and climate changes. In this case, the SOC change is a result of changing practice as well as changing climate. The green line represents a situation in which the farming practices change from current to improved but the climate is the same as in the model initialization phase.

## 13. Discussion

In this discussion chapter we compile the information obtained in the literature survey and the findings of the supportive examples of different types of carbon balance calculations.

### 13.1. Direct measurements

Methods based on felling and weighing of trees can't be used otherwise than in experimental work aiming to produce transfer functions to quantify the carbon content of trees via indirect, like diameter and height, measurements. This means that transfer functions based on non-destructive measurements (diameter at breast height, height) needs to be applied in estimating the stand biomass. The advantage of this is that we can get an estimate of all tree compartments (above and below ground) at the same time. More experimental work is needed to produce spatially representative equations for all tree species. When the inventories are focusing on other types of vegetation than trees, sample based dry-weight measurements could be useful especially for measurements of carbon storages and storage changes of the above ground biomass. Below ground biomass measurements are laborious and the accuracy of them in practical conditions can be low due to the difficulty to collect all roots. The sampling needs to be planned so that the spatial variation can be estimated. Due to the annual variation the date of the sampling possibly can't be named based on a calendar date but needs to be decided based on local weather conditions. If inventories based on sampling were used in a large scale, it would enable the use of specialized labor to decrease the random variation in the estimates.

The large carbon stocks compared to small annual stock changes together with large spatial variation even in a small scale reduces the applicability of repeated measurements as a basis for estimating carbon accumulation into soil on a practical scale. However, we need to take care that in future we have measured results from a wide range of long-term experimental

series of carbon accumulation into soil in order to reduce substantially the knowledge gap related to the changes in soil carbon stocks in versatile climatic conditions, plant cover and soil types.

## 13.2. Spectral methods

Studies to use proximal soil sensors for in situ applications that could estimate SOC in real time and on larger spatial scales have been carried out. At the moment the conclusions are that reasonable price equipment are not accurate enough and the calibration and data processing are tedious. More research needs to be done in terms of selecting the proper spectral range of the sensor, the pre-processing methods, and the calibration techniques.

Remote sensing does not give accurate enough estimations of the changes in carbon storages as such and especially on stand level. The method can be used on general monitoring of vegetation, but we do not know the time needed to increase the accuracy of this method to be applied in the carbon market context.

## 13.3. Gas flux methods

Eddy covariance tower and the measurement equipment are expensive, and because of the expertise of the labor needed to maintain the system and analyze the results are high, also the salary costs are high. For these reasons it is not possible to apply EC widely as a basis for carbon markets of single farms. However, the method can be used to produce focused, high accuracy information of gas exchange and the factors affecting on that in selected experimental sites, representing different types of ecosystems. Flux information could be used for calibrating carbon stock changes quantified by other means. If EC methods were to be used e.g. in model validation, different land use types should be covered in different types of environments over long periods of time.

Gas flux chambers are an effective tool to monitor the short time variation in carbon fluxes related to vegetation, soil or their combination. Measured gas flux data can be applied in estimating the values obtained with other measurements and modeling, but the method can not to be used in producing carbon balances in large scale.

## 13.4. Modeling methods

Simulation of carbon stocks with different models is a widely used approach in estimating carbon stocks and stock changes. The availability of models depends on the model structure and the data available for model calibration. Model simulations are usually performed to provide information about how different practises affect the soil carbon and biomass accumulation. However, the experimental evidence to calibrate especially models of soil component is still too few. The structure and the level of details included in different models vary largely. There are several models to be selected from, but there are no general rules how the selection between the models should be done. One essential principal for the selection is the general applicability of the model.

In the practical use the estimation method needs to be cost efficient and the repeatability and transparency of the system needs to be high. Because the role of soil carbon stocks is large in the ecosystem carbon balance, but the measurement of changes in the soil carbon stocks is difficult, application of models seems to be the way to go forward even if the shortcomings of them are still substantial.

On the level of reporting systems, the level of accuracy of the carbon credits calculated with models needs to be agreed. One of the challenges to overcome is to decide how the areal differences in vegetation species composition, soil properties and climate are taken into account in such calculations. The definition of the required level of accuracy of field measurement methods needs also be done.

## 13.5. Concluding remarks

For annual plants there are available reasonably easily applicable methods to measure the above ground biomass accumulation and its spatial variation directly by weighing. For perennials like trees only nondestructive methods based on tree diameter and height can be used. The availability of such models is quite good. The challenges in obtaining data from below ground processes, both root production and decomposition of dead organic material are large. The direct methods to estimate the changes in soil carbon stocks are tedious and inaccurate. Due to this, it is probable that only modeled estimates of changes in soil carbon stocks will be available also in future. The structures of the models have to be considered thoroughly, so that the performance of them in varying environment can be evaluated in advance. If several types of models are going to be applied, their behavior in different conditions needs to be compared.

# 14. References

- Angelopoulou, T., Tziolas, N., Balafoutis, A., Zalidis, G. & Bochtis, D. 2019, Remote Sensing Techniques for Soil Organic Carbon Estimation: A Review, Remote sensing (Basel, Switzerland), **11**(6), s. 676. DOI 10.3390/rs11060676.
- Asner, G.P., Mascaro, J., Anderson, C., Knapp, D.E., Martin, R.E., Kennedy-Bowdoin, T., van Breugel, M., Davies, S., Hall, J.S., Muller-Landau, H.C., Potvin, C., Sousa, W., Wright, J. & Bermingham, E. 2013, High-fidelity national carbon mapping for resource management and REDD, Carbon balance and management, **8**(1), s. 7. DOI 10.1186/1750-0680-8-7.
- Aubinet, M., Papale, D. & Vesala, T. 2012, Eddy covariance : a practical guide to measurement and data analysis, Springer, Dordrecht ;.
- Baldocchi, D. 2003, Assessing the eddy covariance technique for evaluating carbon dioxide exchange rates of ecosystems: past, present and future, Global Change Biology, **9**(4), s. 479-492. DOI 10.1046/j.1365-2486.2003.00629.x.
- Balser, T.C. & Liang, C. 2011, Microbial production of recalcitrant organic matter in global soils: implications for productivity and climate policy, Nature reviews.Microbiology; Nat Rev Microbiol, **9**(1), s. 75. DOI 10.1038/nrmicro2386-cl.
- Battle-Aguilar, J., Battle-Aguilar, J., Brovelli, A., Brovelli, A., Porporato, A., Porporato, A., Barry, D. & Barry, D. 2011, Modelling soil carbon and nitrogen cycles during land use change. A review, Agronomy for Sustainable Development, **31**(2), s. 251-274. DOI 10.1051/agro/2010007.
- Beets, P., Robertson, K., Ford-Robertson, Gordon, J. and Maclaren, J. 1999. Description and validation of C change: a model for simulating carbon content in managed Pinus Radiata stands. New Zeland Journal of Forestry Science 29:409-427.
- Bellocchi, G., Rivington, M., Donatelli, M. & Matthews, K. 2010, Validation of biophysical models: issues and methodologies. A review, Agronomy for sustainable development, **30**(1), s. 109-130. DOI 10.1051/agro/2009001.
- Berenguer, E., Gardner, T.A., Ferreira, J., Aragão, L.E.O., Camargo, P.B., Cerri, C.E., Durigan, M., Oliveira Junior, R., Vieira, I.C.G. & Barlow, J. 2015, Developing Cost-Effective Field Assessments of Carbon Stocks in Human-Modified Tropical Forests, PloS one; PloS One, **10**(8), s. e0133139. DOI 10.1371/journal.pone.0133139.



- Fassnacht, F.E., Hartig, F., Latifi, H., Berger, C., Hernández, J., Corvalán, P. & Koch, B. 2014, Importance of sample size, data type and prediction method for remote sensing-based estimations of aboveground forest biomass, *Remote sensing of environment*, **154**, s. 102-114. DOI 10.1016/j.rse.2014.07.028.
- Feldpausch, T.R., Lloyd, J., Lewis, S.L., Brienen, R.J.W., Gloor, M., Monteagudo Mendoza, A., Lopez-Gonzalez, G., Banin, L., Abu Salim, K., Affum-Baffoe, K., Alexiades, M., Almeida, S., Amaral, I., Andrade, A., Aragão, L.E.O., Araujo Murakami, A., Arets, E. J. M. M., Arroyo, L., Aymard C, G.A., Baker, T.R., Bánki, O., S., Berry, N.J., Cardozo, N., Chave, J., Comiskey, J.A., Alvarez, E., de Oliveira, A., Di Fiore, A., Djagbletey, G., Domingues, T.F., Erwin, T.L., Fearnside, P.M., França, M., B., Freitas, M.A., Higuchi, N., Iida, Y., Jiménez, E., Kassim, A.R., Killeen, T.J., Laurance, W.F., Lovett, J.C., Malhi, Y., Marimon, B.S., Marimon-Junior, B., Lenza, E., Marshall, A.R., Mendoza, C., Metcalfe, D.J., Mitchard, E.T.A., Neill, D.A., Nelson, B.W., Nilus, R., Nogueira, E.M., Parada, A., Peh, K.S., Pena Cruz, A., Peñuela, M., C., Pitman, N.C.A., Prieto, A., Quesada, C.A., Ramírez, F., Ramírez-Angulo, H., Reitsma, J.M., Rudas, A., Saiz, G., Salomão, R., P., Schwarz, M., Silva, N., Silva-Espejo, J., Silveira, M., Sonké, B., Stropp, J., Taedoumg, H.E., Tan, S., ter Steege, H., Terborgh, J., Torello-Raventos, M., van der Heijden, G. M., F., Vásquez, R., Vilanova, E., Vos, V.A., White, L., Willcock, S., Woell, H. & Phillips, O.L. 2012, Tree height integrated into pantropical forest biomass estimates, *Biogeosciences*, **9**(8), s. 3381-3403. DOI 10.5194/bg-9-3381-2012.
- Flechard CR, Ibrom, A., Skiba, U.M., de Vries, W., van Oijen, M., Cameron, D.R., Dise, N.B., Korhonen, J.F.J., Buchmann, N., Legout, A., Simpson, D., Sanz, M.J., Aubinet, M., Loustau, D., Montagnani, L., Neiryneck, J., Janssens, I.A., Pihlatie, M., Kiese, R., Siemens, J., Francez, A., Augustin, J., Varlagin, A., Olejnik, J., Juszczak, R., Aurela, M., Berveiller, D., Chojnicki, B.H., Dammgen, U., Delpierre, N., Djuricic, V., Drewer, J., Dufrene, E., Eugster, W., Fauvel, Y., Fowler, D., Frumau, A., Granier, A., Gross, P., Hamon, Y., Helfter, C., Hensen, A., Horvath, L., Kitzler, B., Kruijt, B., Kutsch, W.L., Lobo-do-Vale, R., Lohila, A., Longdoz, B., Marek, M., Matteucci, G., Mitasinkova, M., Moreaux, V., Neftel, A., Ourcival, J., Pilegaard, K., Pita, G., Sanz, F., Schjoerring, J.K., Sebastia, M., Tang, Y.S., Uggerud, H., Urbaniak, M., van Dijk, N., Vesala, T., Vidic, S., Vincke, C., Weidinger, T., Zechmeister-Boltenstern, S., Butterbach-Bah, K., Nemitz, E. & Sutton, M.A.: Carbon-nitrogen interactions in European forests and semi-natural vegetation - Part 1 : Fluxes and budgets of carbon, nitrogen and greenhouse gases from ecosystem monitoring and modelling., 2020.

- Freibauer, A., Rounsevell, M.D.A., Smith, P. & Verhagen, J. 2004, Carbon sequestration in the agricultural soils of Europe, *Geoderma*, **122**(1), s. 1-23. DOI 10.1016/j.geoderma.2004.01.021.
- García-Oliva, F. & Masera, O. 2004, Assessment and Measurement Issues Related to Soil Carbon Sequestration in Land-Use, Land-Use Change, and Forestry (LULUCF) Projects under the Kyoto Protocol, *Climatic Change*, **65**(3), s. 347-364. DOI 10.1023/B:CLIM.0000038211.84327.d9.
- Gehl, R.J. & Rice, C.W. 2006, Emerging technologies for in situ measurement of soil carbon, *Climatic Change*, **80**(1-2), s. 43-54. DOI 10.1007/s10584-006-9150-2.
- Goetz, S.J., Baccini, A., Laporte, N.T., Johns, T., Walker, W., Kellndorfer, J., Houghton, R.A. & Sun, M. 2009, Mapping and monitoring carbon stocks with satellite observations: a comparison of methods, *Carbon balance and management*, **4**(1), s. 2. DOI 10.1186/1750-0680-4-2.
- Goidts, E., van Wesemael, B. & Crucifix, M. 2009a, Magnitude and sources of uncertainties in soil organic carbon (SOC) stock assessments at various scales, *European Journal of Soil Science*, **60**(5), s. 723-739. DOI 10.1111/j.1365-2389.2009.01157.x.
- Gold Standard. About us, Vision and Mission. <https://www.goldstandard.org/about-us/vision-and-mission>. Accessed on 12 October 2020.
- Gold Standard. Soil Organic Carbon Framework Methodology. 2020. <https://globalgoals.goldstandard.org/402-luf-agr-fm-soil-organic-carbon-framework-methodology/> Accessed on 12 October 2020.
- Grimault, J., Bellassen, V. and Shishlov, I. 2018. Key elements and challenges in monitoring, certifying and financing forestry carbon projects. Climate brief N58. I4CE – Institute of Climate Economics.
- Gross, A., Hook, L., and Powley, T. Financial news. London DECEMBER 13 2019. <https://www.ft.com/content/7e4665a2-1776-11ea-8d73-6303645ac406>. Accessed on 7 October 2019.
- Hall, F.G., Bergen, K., Blair, J.B., Dubayah, R., Houghton, R., Hurtt, G., Kellndorfer, J., Lefsky, M., Ranson, J., Saatchi, S., Shugart, H.H. & Wickland, D. 2011, Characterizing 3D vegetation structure from space: Mission requirements, *Remote Sensing of Environment*, **115**(11), s. 2753-2775. DOI 10.1016/j.rse.2011.01.024.
- Hame, T., Salli, A., Andersson, K. & Lohi, A. 1997, A new methodology for the estimation of biomass of coniferdominated boreal forest using NOAA AVHRR

data, *International Journal of Remote Sensing*, **18**(15), s. 3211-3243. DOI 10.1080/014311697217053.

Han, S.H. & Park, B.B. 2020, Comparison of Allometric Equation and Destructive Measurement of Carbon Storage of Naturally Regenerated Understory in a *Pinus rigida* Plantation in South Korea, *Forests*, **11**(4), s. 425. DOI 10.3390/f11040425.

Harden, J.W., Hugelius, G., Ahlström, A., Blankinship, J.C., Bond-Lamberty, B., Lawrence, C.R., Loisel, J., Malhotra, A., Jackson, R.B., Ogle, S., Phillips, C., Ryals, R., Todd-Brown, K., Vargas, R., Vergara, S.E., Cotrufo, M.F., Keiluweit, M., Heckman, K.A., Crow, S.E., Silver, W.L., DeLonge, M. & Nave, L.E. 2017, Networking our science to characterize the state, vulnerabilities, and management opportunities of soil organic matter, *Global Change Biology; Glob Chang Biol*, **24**(2), s. e705-e718. DOI 10.1111/gcb.13896.

Harmon, M. & Sexton, J. 1996. Guidelines for measurements of woody detritus in forest ecosystem. Publication No. 20 U.S. LTER Network Office: University of Washington, Seattle, WA, USA. 73 p.

Hbirkou, C., Pätzold, S., Mahlein, A. & Welp, G. 2012, Airborne Birdsey, R., Angeles-Perez, G., Kurz, W.A., Lister, A., Olguin, M., Pan, Y., Wayson, C., Wilson, B. & Johnson, K. 2013, Approaches to monitoring changes in carbon stocks for REDD, *Carbon management*, **4**(5), s. 519-537. DOI 10.4155/cmt.13.49.

Bispo, A., Andersen, L., Angers, D.A., Bernoux, M., Brossard, M., Cécillon, L., Comans, R.N.J., Harmsen, J., Jonassen, K., Lamé, F., Lhuillery, C., Maly, S., Martin, E., Mcelnea, A.E., Sakai, H., Watabe, Y. & Eglin, T.K. 2017, Accounting for Carbon Stocks in Soils and Measuring GHGs Emission Fluxes from Soils: Do We Have the Necessary Standards? *Frontiers in environmental science*, **5**(JUL). DOI 10.3389/fenvs.2017.00041.

Blakemore, L. C., Searle, P. L., Daly, B. K., New Zealand., & New Zealand. 1987. Methods for chemical analysis of soils. Lower Hutt, N.Z: NZ Soil Bureau, Dept. of Scientific and Industrial Research.

Boyero, L., Pearson, R.G., Gessner, M.O., Barmuta, L.A., Ferreira, V., Graça, M.A., S., Dudgeon, D., Boulton, A.J., Callisto, M., Chauvet, E., Helson, J.E., Bruder, A., Albariño, R., J., Yule, C.M., Arunachalam, M., Davies, J.N., Figueroa, R., Flecker, A.S., Ramírez, A., Death, R.G., Iwata, T., Mathooko, J.M., Mathuriau, C., Gonçalves, J., F., Moretti, M.S., Jinggut, T., Lamothe, S., M'Erimba, C., Ratnarajah, L., Schindler, M.H., Castela, J., Buria, L.M., Cornejo, A., Villanueva, V.D. & West, D.C. 2011, A global experiment suggests

climate warming will not accelerate litter decomposition in streams but might reduce carbon sequestration, *Ecology Letters*; *Ecol Lett*, **14**(3), s. 289-294. DOI 10.1111/j.1461-0248.2010.01578.x.

Brown, D.G. 2013, *Land use and the carbon cycle advances in integrated science, management, and policy*, Cambridge University Press, Cambridge ;

Brown, S. 1997. *Estimating Biomass and Biomass Change of Tropical Forests: A Primer*. Rome: FAO Forestry Paper 134.

Byrne, K.A., Kiely, G. & Leahy, P. 2007, Carbon sequestration determined using farm scale carbon balance and eddy covariance, *Agriculture, Ecosystems & Environment*, **121**(4), s. 357-364. DOI 10.1016/j.agee.2006.11.015.

Cairns, M.A., Brown, S., Helmer, E.H. & Baumgardner, G.A. 1997, Root biomass allocation in the world's upland forests, *Oecologia*; *Oecologia*, **111**(1), s. 1-11. DOI 10.1007/s004420050201.

Campbell, E.E. & Paustian, K. 2015, Current developments in soil organic matter modeling and the expansion of model applications: a review, *Environmental research letters*, **10**(12), s. 123004. DOI 10.1088/1748-9326/10/12/123004.

Canada Centre for Remote Sensing. 2019. *Fundamentals of Remote Sensing*. A Canada Centre for Remote Sensing Remote Sensing Tutorial. <https://www.nrcan.gc.ca/maps-tools-publications/satellite-imagery-air-photos/tutorial-fundamentals-remote-sensing/9309> . 8.6.2019. Accessed on 12 October 2020.

Carcia-Shanzes, F., Galvez-Sola, L., Martinez-Nicolas, J., Muelas-Dominigo R. and Nieves, M. 2017. *Using Near-Infrared Spectroscopy in Agricultural Systems*.

Chave, J., Condit, R., Aguilar, S., Hernandez, A., Lao, S. & Perez, R. 2004, Error propagation and scaling for tropical forest biomass estimates, *Philosophical transactions. Biological sciences*; *Philos Trans R Soc Lond B Biol Sci*, **359**(1443), s. 409-420. DOI 10.1098/rstb.2003.1425.

Chen, Y., Chen, Y., Yu, J., Yu, J., Su, X., Su, X., Luo, G. & Luo, G. 2015, Path Planning for Multi-UAV Formation, *Journal of intelligent & robotic systems*, **77**(1), s. 229-246. DOI 10.1007/s10846-014-0077-y.

Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, M. Heimann, C. Jones, C. Le Quéré, R.B. Myneni, S. Piao and P. Thornton, 2013: *Carbon and Other Biogeochemical Cycles*. In: *Climate Change 2013: The Physical*

Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Coleman, K. & Jenkinson, D. 2014, RothC – A model for the turnover of carbon in soil: Model description and users guide (windows version) [Kotisivu: Rothamsted Research]. Saataavilla: <https://search.datacite.org/works/10.23637/model-manual-rothc-win-version-2014>.

Coleman K., Jenkinson D.S. 1996. RothC-26.3 - A Model for the turnover of carbon in soil. In: Powlson D.S., Smith P., Smith J.U. (eds) Evaluation of Soil Organic Matter Models. NATO ASI Series (Series I: Global Environmental Change), vol 38. Springer, Berlin, Heidelberg. [https://doi.org/10.1007/978-3-642-61094-3\\_17](https://doi.org/10.1007/978-3-642-61094-3_17).

Cremers, D.A., Ebinger, M.H., Breshears, D.D., Unkefer, P.J., Kammerdiener, S.A., Ferris, M.J., Catlett, K.M. & Brown, J.R. 2001a, Measuring Total Soil Carbon with Laser-Induced Breakdown Spectroscopy (LIBS), *Journal of Environmental Quality*, **30**(6), s. 2202-2206. DOI 10.2134/jeq2001.2202.

Davidson, E.A. & Janssens, I.A. 2006, Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, *Nature* (London); *Nature*, **440**(7081), s. 165-173. DOI 10.1038/nature04514.

Davis, M., Alves, B., Karlen, D., Kline, K., Galdos, M. & Abulebdeh, D. 2017, Review of Soil Organic Carbon Measurement Protocols: A US and Brazil Comparison and Recommendation, *Sustainability* (Basel, Switzerland), **10**(2), s. 53. DOI 10.3390/su10010053.

Dirzo, R. & Raven, P.H. 2003, GLOBAL STATE OF BIODIVERSITY AND LOSS, *Annual review of environment and resources*, **28**(1), s. 137-167. DOI 10.1146/annurev.energy.28.050302.105532.

Disney, M.I., Boni Vicari, M., Burt, A., Calders, K., Lewis, S.L., Raunonen, P. & Wilkes, P. 2018, Weighing trees with lasers: advances, challenges and opportunities, *Interface focus*; *Interface Focus*, **8**(2), s. 20170048. DOI 10.1098/rsfs.2017.0048.

Duncanson, L., Rourke, O. & Dubayah, R. 2015, Small Sample Sizes Yield Biased Allometric Equations in Temperate Forests, *Scientific reports*; *Sci Rep*, **5**(1), s. 17153. DOI 10.1038/srep17153.

- Elevitch, C., Mazaroli, D. & Ragone, D. 2018, Agroforestry Standards for Regenerative Agriculture, Sustainability (Basel, Switzerland), **10**(9), s. 3337. DOI 10.3390/su10093337.
- Elofsson, K. & Gren, I. 2018, Cost-efficient climate policies for interdependent carbon pools, Environmental modelling & software : with environment data news, **101**, s. 86-101. DOI 10.1016/j.envsoft.2017.12.006.
- England, J.R. & Viscarra Rossel, R.A. 2018, Proximal sensing for soil carbon accounting, Soil, **4**(2), s. 101-122. DOI 10.5194/soil-4-101-2018.
- European Commission. EU climate action and the European Green Deal. [https://ec.europa.eu/clima/policies/eu-climate-action\\_en](https://ec.europa.eu/clima/policies/eu-climate-action_en). Accessed on 11 November 2020.
- European Commission. European Climate Law. [https://ec.europa.eu/clima/policies/eu-climate-action/law\\_en](https://ec.europa.eu/clima/policies/eu-climate-action/law_en) Accessed on 11 November 2020.
- Everaldo de Carvalho Conceição Telles, Plínio Barbosa de Camargo, Luiz A. Martinelli, Susan E. Trumbore, Enir Salazar da Costa, Joaquim Santos, Niro Higuchi & Raimundo Cosme Oliveira Jr 2003, Influence of soil texture on carbon dynamics and storage potential in tropical forest soils of Amazonia, Global biogeochemical cycles, **17**(2), s. 1040-n/a. DOI 10.1029/2002GB001953.
- FAO. 2019. Measuring and modelling soil carbon stocks and stock changes in livestock production systems: Guidelines for assessment (Version 1). Livestock Environmental Assessment and Performance (LEAP) Partnership. Rome, FAO. 170 pp.
- hyperspectral imaging of spatial soil organic carbon heterogeneity at the field-scale, Geoderma, **175-176**, s. 21-28. DOI 10.1016/j.geoderma.2012.01.017.
- Heikkinen J: Carbon storage of Finnish agricultural mineral soils and its long-term change. Helsingin yliopisto: , 2016.
- Heikkinen, J., Kurganova, I., Lopes de Gerenyu, V., Palosuo, T. & Regina, K. 2014a, Changes in soil carbon stock after cropland conversion to grassland in Russian temperate zone: measurements versus model simulation, Nutrient Cycling in Agroecosystems, **98**(1), s. 97-106. DOI 10.1007/s10705-014-9599-8.
- Hobley, E. and Willgoose, G. 2010. Measuring soil organic carbon stocks-issues and consideration. In: Proceedings of the 19<sup>th</sup> World Congress of Soil Science: Soil

solutions for a changing world, Brisbane, Australia, 1-6 August 2010. Symposium 1.5.1 Quantitative monitoring of soil change 2010 pp.62-65 ref.21

Houghton, R.A. 2003, Why are estimates of the terrestrial carbon balance so different? *Global Change Biology*, **9**(4), s. 500-509. DOI 10.1046/j.1365-2486.2003.00620.x.

Houghton, R.A., Hall, F. & Goetz, S.J. 2009, Importance of biomass in the global carbon cycle, *Journal of Geophysical Research: Biogeosciences*, **114**, s. G00E03-n/a. DOI 10.1029/2009JG000935.

Husch, B. & Beers, T. W. & Kershaw, Jr, J. A. 2003. *Forest Mensuration*. 4. painos. Hoboken, New Jersey, USA: John Wiley & Sons, Inc. 443 s.

Hynynen, J.; Salminen, H.; Ahtikoski, A.; Huuskonen, S.; Ojansuu, R.; Siipilehto, J.; Lehtonen, M.; Rummukainen, A.; Kojola, S.; Eerikäinen, K. Scenario Analysis for the Biomass Supply Potential and the Future Development of Finnish Forest Resources. Working Papers of the Finnish Forest Research Institute. 2014, 302, 106.

Iizuka, K. & Tateishi, R. 2015, Estimation of CO<sub>2</sub> Sequestration by the Forests in Japan by Discriminating Precise Tree Age Category using Remote Sensing Techniques, *Remote sensing (Basel, Switzerland)*, **7**(11), s. 15082-15113. DOI 10.3390/rs71115082.

IPCC Guidelines for National Greenhouse Gas Inventories. 2006. Volume 4: Agriculture, Forestry and Other Land Use. [https://www.ipccnggip.iges.or.jp/public/2006gl/pdf/4\\_Volume4/V4\\_01\\_Ch1\\_Introduction.pdf](https://www.ipccnggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_01_Ch1_Introduction.pdf) ipcc 2006

IPCC. 2003. Good practice guidance for land use, land-use change and forestry. Penman, J., Gytarsky, M., Hiraishi, T., Krug, T., Kruger, D., Pipatti, R., Buen-día, L., Miwa, K., Ngara, T., Tanabe, K. & Wagner, F. (Eds.) Intergovernmental Panel on Climate Change. IGES, Hayama, Japan. 593 pp.

Issa, S., Dahy, B., Ksiksi, T. & Saleous, N. 2020, A Review of Terrestrial Carbon Assessment Methods Using Geo-Spatial Technologies with Emphasis on Arid Lands, *Remote sensing (Basel, Switzerland)*, **12**(12), s. 2008. DOI 10.3390/rs12122008.

Izaurralde, R.C., Rice, C.W., Wielopolski, L., Ebinger, M.H., Reeves, J., James B, Thomson, A.M., Harris, R., Francis, B., Mitra, S., Rappaport, A.G., Etchevers, J.D., Sayre, K.D., Govaerts, B. & McCarty, G.W. 2013, Evaluation of Three Field-Based Methods for Quantifying Soil Carbon, *PloS one*, **8**(1), s. e55560. DOI 10.1371/journal.pone.0055560.



- Jackson, R.B., Lajtha, K., Crow, S.E., Hugelius, G., Kramer, M.G. & Piñeiro, G. 2017, The Ecology of Soil Carbon: Pools, Vulnerabilities, and Biotic and Abiotic Controls, *Annual review of ecology, evolution, and systematics*, **48**(1), s. 419-445. DOI 10.1146/annurev-ecolsys-112414-054234.
- Jackson, R.B., Mooney, H.A. & Schulze, E.-D. 1997, A global budget for fine root biomass, surface area, and nutrient contents, *Proceedings of the National Academy of Sciences – PNAS; Proc Natl Acad Sci U S A*, **94**(14), s. 7362-7366. DOI 10.1073/pnas.94.14.7362.
- Janzen, H.H. 2004, Carbon cycling in earth systems—a soil science perspective, *Agriculture, Ecosystems & Environment*, **104**(3), s. 399-417. DOI 10.1016/j.agee.2004.01.040.
- Jari Liski, Mikko Tuomi & Jussi Rasinmäki Yasso07 user-interface manual.
- Jobbagy, E.G. & Jackson, R.B. 2000, The Vertical Distribution of Soil Organic Carbon and Its Relation to Climate and Vegetation, *Ecological applications*, **10**(2), s. 423. DOI 10.2307/2641104.
- Jones & Fallon 2009 Sources of uncertainty in global modelling of future SOC storage Book *Uncertainties in environmental modelling a*,
- Kallenbach, C.M., Frey, S.D. & Grandy, A.S. 2016, Direct evidence for microbial-derived soil organic matter formation and its ecophysiological controls, *Nature communications; Nat Commun*, **7**(1), s. 13630. DOI 10.1038/ncomms13630.
- Karhu, K., Gärdenäs, A., I., Heikkinen, J., Vanhala, P., Tuomi, M. & Liski, J. 2012, Impacts of organic amendments on carbon stocks of an agricultural soil — Comparison of model-simulations to measurements, *Geoderma*, **189-190**, s. 606-616. DOI 10.1016/j.geoderma.2012.06.007.
- Kate M. Buckeridge, Kelly E. Mason, Niall P. McNamara, Nick Ostle, Jeremy Puissant, Tim Goodall, Robert I. Griffiths, Andrew W. Stott & Jeanette Whitaker: Environmental and microbial controls on microbial necromass recycling, an important precursor for soil carbon stabilization. SUKUNIMET VÄÄRIN
- Kätterer, T. 2014, C-TOOL: A simple model for simulating whole-profile carbon storage in temperate agricultural soils.
- Kätterer, T., Reichstein, M., Andrén, O. & Lomander, A. 1998, Temperature dependence of organic matter decomposition: a critical review using literature data analyzed with different models, *Biology and Fertility of Soils*, **27**(3), s. 258-262. DOI 10.1007/s003740050430.

- Kelleher, B.P. & Simpson, A.J. 2006, Humic Substances in Soils: Are They Really Chemically Distinct? *Environmental science & technology; Environ.Sci.Technol*, **40**(15), s. 4605-4611. DOI 10.1021/es0608085.
- Keller, M., Palace, M. & Hurtt, G. 2001, Biomass estimation in the Tapajos National Forest, Brazil: Examination of sampling and allometric uncertainties, *Forest Ecology and Management*, **154**(3), s. 371-382. DOI 10.1016/S0378-1127(01)00509-6.
- Killham, K. & Foster, R. 1994, *Soil ecology*, Cambridge University Press, Cambridge.
- King RL: A challenge for high spatial, spectral, and temporal resolution data fusion. *IEEE: 2602-2604 vol.6*, 2000.
- Knoke, T. 2013, Uncertainties and REDD+: Implications of applying the conservativeness principle to carbon stock estimates, *Climatic Change*, **119**(2), s. 261-267. DOI 10.1007/s10584-012-0662-7.
- Köhl, M., Neupane, P.R. & Mundhenk, P. 2020, REDD+ measurement, reporting and verification – A cost trap? Implications for financing REDD+MRV costs by result-based payments, *Ecological Economics*, **168**, s. 106513. DOI 10.1016/j.ecolecon.2019.106513.
- Koolhaas, R. 2014, *Fundamentals*, 1. ed. p., Marsilio, Venezia.
- Körschens, M. 2005, Global and regional importance of long-term field experiments, *Archiv für Acker- und Pflanzenbau und Bodenkunde*, **51**(2), s. 111-117. DOI 10.1080/03650340400026677.
- Kuzyakov, Y. 2010, Priming effects: Interactions between living and dead organic matter, *Soil biology & biochemistry*, **42**(9), s. 1363-1371. DOI 10.1016/j.soilbio.2010.04.003.
- Lasslop, G., Reichstein, M., Kattge, J. & Papale, D. 2008, Influences of observation errors in eddy flux data on inverse model parameter estimation, *Biogeosciences*, **5**(5), s. 1311-1324. DOI 10.5194/bg-5-1311-2008.
- LeBauer, D.S. & Treseder, K.K. 2008, NITROGEN LIMITATION OF NET PRIMARY PRODUCTIVITY IN TERRESTRIAL ECOSYSTEMS IS GLOBALLY DISTRIBUTED, *Ecology (Durham); Ecology*, **89**(2), s. 371-379. DOI 10.1890/06-2057.1.
- Lindner, M., Lindner, M., Karjalainen, T. & Karjalainen, T. 2007, Carbon inventory methods and carbon mitigation potentials of forests in Europe: a short review of recent progress, *European Journal of Forest Research*, **126**(2), s. 149-156. DOI 10.1007/s10342-006-0161-3.

- Liski, J., Lehtonen, A., Palosuo, T., Peltoniemi, M., Eggers, T., Muukkonen, P. & Mäkipää, R. 2006, Carbon accumulation in Finland's forests 1922–2004 – an estimate obtained by combination of forest inventory data with modelling of biomass, litter and soil, *Annals of forest science.*, **63**(7), s. 687-697. DOI 10.1051/forest:2006049.
- Liski, J., Palosuo, T., Peltoniemi, M. & Sievänen, R. 2005, Carbon and decomposition model Yasso for forest soils, *Ecological modelling*, **189**(1-2), s. 168-182. DOI 10.1016/j.ecolmodel.2005.03.005.
- Liu B & Liu, W.: Fusion of Multi-spectral Image Using Non-separable Additive Wavelets for High Spatial Resolution Enhancement. Springer Berlin Heidelberg: 90-99, Berlin, Heidelberg.
- Liu, D.L., Chan, K.Y., Conyers, M.K., Li, G. & Poile, G.J. 2011, Simulation of soil organic carbon dynamics under different pasture managements using the RothC carbon model, *Geoderma*, **165**(1), s. 69-77. DOI 10.1016/j.geoderma.2011.07.005.
- Lorenz, K. & Lal, R. 2018, *Carbon Sequestration in Agricultural Ecosystems*, Springer International Publishing, Cham.
- Lu, D. 2007, The potential and challenge of remote sensing-based biomass estimation, *International journal of remote sensing*, **27**(7), s. 1297-1328. DOI 10.1080/01431160500486732.
- Lukac, M. & Godbold, D.L. 2011, *Soil ecology in northern forests a belowground view of a changing world*, Cambridge University Press, Cambridge ;.
- Luo, Z., Feng, W., Luo, Y., Baldock, J. & Wang, E. 2017, Soil organic carbon dynamics jointly controlled by climate, carbon inputs, soil properties and soil carbon fractions, *Global Change Biology; Glob Chang Biol*, **23**(10), s. 4430-4439. DOI 10.1111/gcb.13767.
- Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S., Grelle, A., Hari, P., Jarvis, P.G., Kolari, P., Kowalski, A.S., Lankreijer, H., Law, B.E., Lindroth, A., Loustau, D., Manca, G., Moncrieff, J.B., Rayment, M., Tedeschi, V., Valentini, R. & Grace, J. 2007, The human footprint in the carbon cycle of temperate and boreal forests, *Nature (London); Nature*, **447**(7146), s. 849-851. DOI 10.1038/nature05847.
- Mäkelä, A. & Valentine, H.T. 2019, *Models of Tree and Stand Dynamics : Theory, Formulation and Application*, Springer International Publishing.

- Mäkipää, R. 1995, Effect of nitrogen input on carbon accumulation of boreal forest soils and ground vegetation, *Forest Ecology and Management*, **79**(3), s. 217-226. DOI 10.1016/0378-1127(95)03601-6.
- Näsi, L., Hänninen, H. & Anttonen, J. 2008a, The costs of monitoring changes in forest soil carbon stocks, *Boreal Environment Research Publishing Board*.
- Mäkipää, R., Karjalainen, T., Pussinen, A. & Kukkola, M. 1998, Effects of nitrogen fertilization on carbon accumulation in boreal forests: Model computations compared with the results of long-term fertilization experiments, *Chemosphere (Oxford)*, **36**(4), s. 1155-1160. DOI 10.1016/S0045-6535(97)10188-6.
- Manzoni, S. & Porporato, A. 2009, Soil carbon and nitrogen mineralization: Theory and models across scales, *Soil biology & biochemistry*, **41**(7), s. 1355-1379. DOI 10.1016/j.soilbio.2009.02.031.
- Mayzelle, M.M., Krusor, M.L., Lajtha, K., Bowden, R.D. & Six, J. 2014, Effects of Detrital Inputs and Roots on Carbon Saturation Deficit of a Temperate Forest Soil, *Soil Science Society of America Journal*, **78**, s. S76-S83. DOI 10.2136/sssaj2013.09.0415nafsc.
- Miltner, A., Bombach, P., Schmidt-Brücken, B. & Kästner, M. 2011, SOM genesis: microbial biomass as a significant source, *Biogeochemistry*, **111**(1-3), s. 41-55. DOI 10.1007/s10533-011-9658-z.
- Minasny, B., Malone, B.P., McBratney, A.B., Angers, D.A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z., Cheng, K., Das, B.S., Field, D.J., Gimona, A., Hedley, C.B., Hong, S.Y., Mandal, B., Marchant, B.P., Martin, M., McConkey, B.G., Mulder, V.L., O'Rourke, S., Richer-de-Forges, A.C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C., Vågen, T., van Wesemael, B. & Winowiecki, L. 2017, Soil carbon 4 per mille, *Geoderma*, **292**, s. 59-86. DOI 10.1016/j.geoderma.2017.01.002.
- Minx, J.C., Lamb, W.F., Callaghan, M.W., Fuss, S., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., de Oliveira Garcia, W., Hartmann, J., Khanna, T., Lenzi, D., Luderer, G., Nemet, G.F., Rogelj, J., Smith, P., Vicente Vicente, J.L., Wilcox, J. & del Mar Zamora Dominguez, Maria 2018, Negative emissions—Part 1: Research landscape and synthesis, *Environmental research letters*, **13**(6), s. 063001. DOI 10.1088/1748-9326/aabf9b.
- Moore, J.R. 2010, Allometric equations to predict the total above-ground biomass of radiata pine trees, *Annals of forest science.*, **67**(8), s. 806. DOI 10.1051/forest/2010042.

- Morvan, X., Saby, N.P.A., Arrouays, D., Le Bas, C., Jones, R.J.A., Verheijen, F.G.A., Bellamy, P.H., Stephens, M. & Kibblewhite, M.G. 2008, Soil monitoring in Europe: A review of existing systems and requirements for harmonisation, *The Science of the total environment*; *Sci Total Environ*, **391**(1), s. 1-12. DOI 10.1016/j.scitotenv.2007.10.046.
- Muukkonen, P. & Heiskanen, J. 2007, Biomass estimation over a large area based on standwise forest inventory data and ASTER and MODIS satellite data: A possibility to verify carbon inventories, *Remote sensing of environment*, **107**(4), s. 617-624. DOI 10.1016/j.rse.2006.10.011.
- Nawar, S., Buddenbaum, H., Hill, J., Kozak, J. & Mouazen, A.M. 2016, Estimating the soil clay content and organic matter by means of different calibration methods of vis-NIR diffuse reflectance spectroscopy, *Soil & tillage research*, **155**, s. 510-522. DOI 10.1016/j.still.2015.07.021.
- Nelson, D.W., and L.E. Sommers. 1996. Total carbon, organic carbon, and organic matter. In: D.L. Sparks, editor, *Methods of soil analysis. Part 3 Chemical methods*. SSSA and ASA, Madison, WI. p. 961-1010
- Ogle, S.M., Breidt, F.J. & Paustian, K. 2006, Bias and variance in model results associated with spatial scaling of measurements for parameterization in regional assessments, *Global Change Biology*, **12**(3), s. 516-523. DOI 10.1111/j.1365-2486.2006.01106.x.
- Ollikainen, M., Lankoski, J. & Lötjönen, S. 2020, Climate change mitigation and agriculture: measures, costs and policies – A literature review, *Agricultural and food science*, **29**(2). DOI 10.23986/afsci.85830.
- Ondrasek, G., Bakić Begić, H., Zovko, M., Filipović, L., Meriño-Gergichevich, C., Savić, R. & Rengel, Z. 2019, Biogeochemistry of soil organic matter in agroecosystems & environmental implications, *The Science of the total environment*; *Sci Total Environ*, **658**, s. 1559-1573. DOI 10.1016/j.scitotenv.2018.12.243.
- Parton, W.J., Hartman, M., Ojima, D. & Schimel, D. 1998, DAYCENT and its land surface submodel: description and testing, *Global and planetary change*, **19**(1-4), s. 35-48. DOI 10.1016/s0921-8181(98)00040-x.
- Parton, W., Schimel, D., Cole, C. & Ojima, D. 1987. Analysis of factors controlling soil organic matter levels in Great Plains grasslands 1. *Soil Science Society of America Journal*, 51:1173-1179. <https://doi.org/10.2136/sssaj.1987.03615.99500.51000.50015x>.
- Paul, E.A. & Paul, E.A. 2007a, *Soil microbiology, ecology, and biochemistry*, 3. p., Academic Press, Amsterdam.

- Paustian, K., Collier, S., Baldock, J., Burgess, R., Creque, J., DeLonge, M., Dungait, J., Ellert, B., Frank, S., Goddard, T., Govaerts, B., Grundy, M., Henning, M., Izaurrealde, R.C., Madaras, M., McConkey, B., Porzig, E., Rice, C., Searle, R., Seavy, N., Skalsky, R., Mulhern, W. & Jahn, M. 2019a, Quantifying carbon for agricultural soil management: from the current status toward a global soil information system, *Carbon management*, **10**(6), s. 567-587. DOI 10.1080/17583004.2019.1633231.
- Paustian, K., Collier, S., Baldock, J., Burgess, R., Creque, J., DeLonge, M., Dungait, J., Ellert, B., Frank, S., Goddard, T., Govaerts, B., Grundy, M., Henning, M., Izaurrealde, R.C., Madaras, M., McConkey, B., Porzig, E., Rice, C., Searle, R., Seavy, N., Skalsky, R., Mulhern, W. & Jahn, M. 2019b, Quantifying carbon for agricultural soil management: from the current status toward a global soil information system, *Carbon management*, **10**(6), s. 567-587. DOI 10.1080/17583004.2019.1633231.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P. & Smith, P. 2016, Climate-smart soils, *Nature (London)*; *Nature*, **532**(7597), s. 49-57. DOI 10.1038/nature17174.
- Pearson TRH: Measurement guidelines for the sequestration of forest carbon / Timothy R.H. Pearson, Sandra L. Brown, Richard A. Birdsey. U.S. Dept. of Agriculture, Forest Service, Northern Research Station: Pennsylvania, 2007.
- Peltoniemi, M., Mäkipää, R., Liski, J. & Tamminen, P. 2004, Changes in soil carbon with stand age – an evaluation of a modelling method with empirical data, *Global Change Biology*, **10**(12), s. 2078-2091. DOI 10.1111/j.1365-2486.2004.00881.x.
- Peltoniemi, M., Mäkipää, R., Liski, J. & Tamminen, P. 2004, Changes in soil carbon with stand age – an evaluation of a modelling method with empirical data, *Global Change Biology*, **10**(12), s. 2078-2091. DOI 10.1111/j.1365-2486.2004.00881.x.
- Peltoniemi, M., Thürig, E., Ogle, S., Palosuo, T., Schrumpf, M., Wutzler, T., Butterbach-Bahl, K., Chertov, O., Komarov, A., Mikhailov, A., Gärdenäs, A., Perry, C., Liski, J., Smith, P. & Mäkipää, R. 2007, Models in country scale carbon accounting of forest soils, *Silva Fennica*, **41**(3). DOI 10.14214/sf.290.
- Peterson, B.L., Hanna, L. & Steiner, J.L. 2019, Reduced soil disturbance: Positive effects on greenhouse gas efflux and soil N losses in winter wheat systems of the southern plains, *Soil & tillage research*, **191**, s. 317-326. DOI 10.1016/j.still.2019.03.020.

- Poeplau, C. & Don, A. 2015, Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-analysis, *Agriculture, Ecosystems & Environment*, **200**, s. 33-41. DOI 10.1016/j.agee.2014.10.024.
- Ponce-Hernandez R, Koohafkan, P. & Antoine, J.: *Assessing Carbon Stocks and Modelling Win-win Scenarios of Carbon Sequestration Through Land-use Changes*. Rome, Italy: Food and Agriculture Organization of the United Nations: , 2004.
- Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P. & Mommer, L. 2012, Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control, *The New phytologist; New Phytol*, **193**(1), s. 30-50. DOI 10.1111/j.1469-8137.2011.03952.x.
- Post, J., Krysanova, V., Suckow, F., Mirschel, W., Rogasik, J. & Merbach, I. 2007, Integrated eco-hydrological modelling of soil organic matter dynamics for the assessment of environmental change impacts in meso- to macro-scale river basins, *Ecological Modelling*, **206**(1), s. 93-109. DOI 10.1016/j.ecolmodel.2007.03.028.
- Posudin, I.I.I. 2014, *Methods of measuring environmental parameters*, Wiley, Hoboken, New Jersey.
- Prasad, V.K., Kant, Y. & Badarinath, K.V.S. 2001, CENTURY ecosystem model application for quantifying vegetation dynamics in shifting cultivation areas: A case study from Rampa Forests, Eastern Ghats (India), *Ecological research*, **16**(3), s. 497-507. DOI 10.1046/j.1440-1703.2001.00412.x.
- Pukkala, T. 2016, Which type of forest management provides most ecosystem services? *Forest ecosystems*, **3**(1), s. 1-16. DOI 10.1186/s40663-016-0068-5.
- Raich, J.W. & Nadelhoffer, K.J. 1989, Belowground Carbon Allocation in Forest Ecosystems: Global Trends, *Ecology (Durham)*, **70**(5), s. 1346-1354. DOI 10.2307/1938194.
- Rantakari, M., Lehtonen, A., Linkosalo, T., Tuomi, M., Tamminen, P., Heikkinen, J., Liski, J., Mäkipää, R., Ilvesniemi, H. & Sievänen, R. 2012, The Yasso07 soil carbon model – Testing against repeated soil carbon inventory, *Forest ecology and management*, **286**, s. 137-147. DOI 10.1016/j.foreco.2012.08.041.
- Rex, F.E., Silva, C.A., Dalla Corte, A.P., Klauberg, C., Mohan, M., Cardil, A., Silva, V.S.d., Almeida, Danilo Roberti Alves de, Garcia, M., Broadbent, E.N., Valbuena, R., Stoddart, J., Merrick, T. & Hudak, A.T. 2020, Comparison of Statistical Modelling Approaches for Estimating Tropical Forest Aboveground Biomass Stock and Reporting Their Changes in Low-Intensity Logging Areas Using Multi-Temporal LiDAR Data, *Remote sensing (Basel, Switzerland)*, **12**(9), s. 1498. DOI 10.3390/rs12091498.



- Richards, G.P. & Evans, D.M.W. 2004, Development of a carbon accounting model (FullCAM Vers. 1.0) for the Australian continent, *Australian forestry*, **67**(4), s. 277-283. DOI 10.1080/00049158.2004.10674947.
- Richards, G.P. & Evans, D.M.W. 2004, Development of a carbon accounting model (FullCAM Vers. 1.0) for the Australian continent, *Australian forestry*, **67**(4), s. 277-283. DOI 10.1080/00049158.2004.10674947.
- Rinne J, Karl, T. & Guenther, A.: Simple, stable, and affordable : Towards long-term ecosystem scale flux measurements of VOCs., 2016.
- Roxburgh, S.H., Paul, K.I., Clifford, D., England, J.R. & Raison, R.J. 2015, Guidelines for constructing allometric models for the prediction of woody biomass: How many individuals to harvest? *Ecosphere*, **6**(3), s. art38-27. DOI 10.1890/ES14-00251.1.
- Saarsalmi, A., Smolander, A., Kukkola, M., Moilanen, M. and Saramäki, J. 2012. 30-Year effects of wood ash and nitrogen fertilization on soil chemical properties, soil microbial processes and stand growth in a Scots pine stand. *Forest Ecology and Management* 278 (2012) 63–70. <http://dx.doi.org/10.1016/j.foreco.2014.04.031>
- Salminen, H.; Lehtonen, M.; Hynynen, J. Reusing Legacy FORTRAN in the MOTTI Growth and Yield Simulator. *Comput. Electron. Agric.* 2005, 49 (1), 103–113. <https://doi.org/10.1016/j.compag.2005.02.005>.
- Sanderman, J., Hengl, T. & Fiske, G.J. 2017, Soil carbon debt of 12,000 years of human land use, *Proceedings of the National Academy of Sciences - PNAS; Proc Natl Acad Sci U S A*, **114**(36), s. 9575-9580. DOI 10.1073/pnas.1706103114.
- Scharlemann, J.P.W., Tanner, E.V.J., Hiederer, R. & Kapos, V. 2014, Global soil carbon: understanding and managing the largest terrestrial carbon pool, *Carbon management*, **5**(1), s. 81-91. DOI 10.4155/cmt.13.77.
- Schimel, D.S., Braswell, B.H., Holland, E.A., McKeown, R., Ojima, D.S., Painter, T.H., Parton, W.J. & Townsend, A.R. 1994, Climatic, edaphic, and biotic controls over storage and turnover of carbon in soils, *Global Biogeochemical Cycles*, **8**(3), s. 279-293. DOI 10.1029/94gb00993.
- Schrumpf, M., Schulze, E.D., Kaiser, K. & Schumacher, J. 2011, How accurately can soil organic carbon stocks and stock changes be quantified by soil inventories? *Biogeosciences*, **8**(5), s. 1193-1212. DOI 10.5194/bg-8-1193-2011.
- Sedjo, R. & Sohngen, B. 2012, Carbon Sequestration in Forests and Soils, *Annual review of resource economics*, **4**(1), s. 127-144. DOI 10.1146/annurev-resource-083110-115941.

- Senapati N, Ghosh, S., Daniel, H. & Rakshit, A.: A Basic Concept on Modelling Soil Organic Carbon. Springer International Publishing: 293-313, Cham, 2014.
- Senapati N., Ghosh S., Daniel H., Rakshit A. (2014) A Basic Concept on Modelling Soil Organic Carbon. In: Basu S., Kumar N. (eds) Modelling and Simulation of Diffusive Processes. Simulation Foundations, Methods and Applications. Springer, Cham. [https://doi-org.libproxy.helsinki.fi/10.1007/978-3-319-05657-9\\_14](https://doi-org.libproxy.helsinki.fi/10.1007/978-3-319-05657-9_14)
- Sinfield, J.V., Fagerman, D. & Colic, O. 2010, Evaluation of sensing technologies for on-the-go detection of macro-nutrients in cultivated soils, Computers and Electronics in Agriculture, **70**(1), s. 1-18. DOI 10.1016/j.compag.2009.09.017.
- Singh, K., Murphy, B.W. & Aubinethant, B.P. 2012, Towards cost-effective estimation of soil carbon stocks at the field scale, Soil research (Collingwood, Vic.), **50**(8), s. 672. DOI 10.1071/SR12119.
- Smith, P. 2004, How long before a change in soil organic carbon can be detected? Global change biology, **10**(11), s. 1878-1883. DOI 10.1111/j.1365-2486.2004.00854.x.
- Smith, P. 2012, Soils and climate change, Current opinion in environmental sustainability, **4**(5), s. 539-544. DOI 10.1016/j.cosust.2012.06.005.
- Smith, P. 2016, Soil carbon sequestration and biochar as negative emission technologies, Global Change Biology; Glob Chang Biol, **22**(3), s. 1315-1324. DOI 10.1111/gcb.13178.
- Smith, P., Lanigan, G., Kutsch, W.L., Buchmann, N., Eugster, W., Aubinet, M., Ceschia, E., Béziat, P., Yeluripati, J.B., Osborne, B., Moors, E.J., Brut, A., Wattenbach, M., Saunders, M. & Jones, M. 2010, Measurements necessary for assessing the net ecosystem carbon budget of croplands, Agriculture, Ecosystems & Environment, **139**(3), s. 302-315. DOI 10.1016/j.agee.2010.04.004.
- Smith, P., Smith, J.U., Powlson, D.S., McGill, W.B., Arah, J.R.M., Chertov, O.G., Coleman, K., Franko, U., Frolking, S., Jenkinson, D.S., Jensen, L.S., Kelly, R.H., Klein-Gunnewiek, H., Komarov, A.S., Li, C., Molina, J.A.E., Mueller, T., Parton, W.J., Thornley, J.H.M. & Whitmore, A.P. 1997, A comparison of the performance of nine soil organic matter models using datasets from seven long-term experiments, Geoderma, **81**(1-2), s. 153-225. DOI 10.1016/s0016-7061(97)00087-6.
- Smith, P., Soussana, J.F., Angers, D., Schipper, L., Chenu, C., Rasse, D.P., Batjes, N.H., Egmond, v., Fenny, McNeill, S., Kuhnert, M., Arias-Navarro, C., Olesen, J.E., Chirinda, N., Fornara, D., Wollenberg, E., Álvaro-Fuentes, J., Sanz-Cobena, A. & Klumpp, K. 2020,

How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal, *Global Change Biology*, **26**(1), s. 219-241. DOI 10.1111/gcb.14815.

Smith, P.E., Anen, O., Brussaard, L., Dangerfield, M., Ekschmitt, K., Lavelle, P. & Tate, K. 1998, Soil biota and global change at the ecosystem level: describing soil biota in mathematical models, *Global Change Biology*, **4**(7), s. 773-784. DOI 10.1046/j.1365-2486.1998.00193.x.

Song, X., Hansen, M.C., Stehman, S.V., Potapov, P.V., Tyukavina, A., Vermote, E.F. & Townshend, J.R. 2018, Global land change from 1982 to 2016, *Nature* (London); *Nature*, **560**(7720), s. 639-643. DOI 10.1038/s41586-018-0411-9.

Statistics Finland. Greenhouse Gas Emissions in Finland 1990-2017. National Inventory Report Under the UNFCCC and the Kyoto Protocol. Statistics Finland, Helsinki, Finland (2019). Available at: <https://unfccc.int/process-and-meetings/transparency-and-reporting/reporting-and-review-under-the-convention/greenhouse-gas-inventories-annex-i-parties/national-inventory-submissions-2019>

Steffen, W., Grinevald, J., Crutzen, P. & McNeill, J. 2011, The Anthropocene: conceptual and historical perspectives, *Philosophical transactions of the Royal Society of London. Series A: Mathematical, physical, and engineering sciences*; *Philos Trans A Math Phys Eng Sci*, **369**(1938), s. 842-867. DOI 10.1098/rsta.2010.0327.

Stenberg, B., Viscarra Rossel, R.,A., Mouazen, A.M. & Wetterlind, J. 2010, Visible and Near Infrared Spectroscopy in Soil Science, *Advances in Agronomy*; *Advances in Agronomy*, **107**, s. 163-215. DOI 10.1016/s0065-2113(10)07005-7.

Stevens, A., van Wesemael, B., Vandenschrick, G., Touré, S. & Tychon, B. 2006, Detection of Carbon Stock Change in Agricultural Soils Using Spectroscopic Techniques, *Soil Science Society of America Journal*, **70**(3), s. 844-850. DOI 10.2136/sssaj2005.0025.

Stockmann, U., Adams, M.A., Crawford, J.W., Field, D.J., Henakaarchchi, N., Jenkins, M., Minasny, B., McBratney, A.B., Courcelles, Vivien de Remy de, Singh, K., Wheeler, I., Abbott, L., Angers, D.A., Baldock, J., Bird, M., Brookes, P.C., Chenu, C., Jastrow, J.D., Lal, R., Lehmann, J., O'Donnell, A.,G., Parton, W.J., Whitehead, D. & Zimmermann, M. 2013, The knowns, known unknowns and unknowns of sequestration of soil organic carbon, *Agriculture, Ecosystems & Environment*, **164**, s. 80-99. DOI 10.1016/j.agee.2012.10.001.

Sullivan, M.J.P., Lewis, S.L., Hubau, W., Qie, L., Baker, T.R., Banin, L.F., Chave, J., Cuni-Sanchez, A., Feldpausch, T.R., Lopez-Gonzalez, G., Arets, E., Ashton, P., Bastin, J., Berry, N.J.,

Bogaert, J., Boot, R., Brearley, F.Q., Brienen, R., Burslem, David F. R. P., de Canniere, C., Chudomelová, M., Dančák, M., Ewango, C., Hédli, R., Lloyd, J., Makana, J., Malhi, Y., Marimon, B.S., Junior, B.H.M., Metali, F., Moore, S., Nagy, L., Vargas, P.N., Pendry, C.A., Ramírez-Angulo, H., Reitsma, J., Rutishauser, E., Salim, K.A., Sonké, B., Sukri, R.S., Sunderland, T., Svátek, M., Umunay, P.M., Martinez, R.V., Vernimmen, R.R.E., Torre, E.V., Vleminckx, J., Vos, V. & Phillips, O.L. 2018, Field methods for sampling tree height for tropical forest biomass estimation, *Methods in ecology and evolution*; *Methods Ecol Evol*, **9**(5), s. 1179-1189. DOI 10.1111/2041-210x.12962.

Sulzman, E.W., Brant, J.B., Bowden, R.D. & Lajtha, K. 2005, Contribution of aboveground litter, belowground litter, and rhizosphere respiration to total soil CO<sub>2</sub> efflux in an old growth coniferous forest, *Biogeochemistry*, **73**(1), s. 231-256. DOI 10.1007/s10533-004-7314-6.

Sun, W. & Liu, X. 2019, Review on carbon storage estimation of forest ecosystem and applications in China, *Forest ecosystems*, **7**(1), s. 1-14. DOI 10.1186/s40663-019-0210-2.

Sun, X., Li, G., Wang, M. & Fan, Z. 2019, Analyzing the Uncertainty of Estimating Forest Aboveground Biomass Using Optical Imagery and Spaceborne LiDAR, *Remote sensing (Basel, Switzerland)*, **11**(6), s. 722. DOI 10.3390/rs11060722.

Sutton Ian 2015, 10.1 Introduction, *teoksessa Process Risk and Reliability Management*, 2nd Edition. p., Elsevier, s. 1-2.

Taiz, L. & Zeiger, E. 2010, *Plant physiology*, 5. p., Sinauer Associates, Sunderland, MA.

To the Carbon Credits (Carbon Farming Initiative—Measurement of Soil Carbon Sequestration in Agricultural Systems) Methodology Determination 2018 Version 1.0 – January 2018: The Supplement.


Toensmeier, E. 2016. *The Carbon Farming Solution. A Global Toolkit of Perennial Crops and Regenerative Agriculture Practices for Climate Change Mitigation and Food Security*. White River Junction, Vermont, USA: Chelsea Green Publishing. 480 s.

Tomppo, E. & Gschwantner, T. & Lawrence, M. & McRoberts, R. E. 2010. *National Forest Inventories. Pathways for Common Reporting*. Springer Science + Business Media. 612 s.

Tomppo, E. 2014, *The multi-source national forest inventory of Finland : methods and results 2011*, Metla, Vantaa.

- Tuomi M, Rasinmäki J, Repo A, Vanhala P, Liski J. 2011. Soil carbon model Yasso07 graphical user interface. *Environ. Model. Softw.* 26(11), 1358-1362.
- Tuomi M, Vanhala P, Karhu K, Fritze H, Liski J. 2008. Heterotrophic soil respiration - comparison of different models describing its temperature dependence. *Ecol. Model.* 211 (1-2), 182-190.
- United nations. What is the Paris Agreement? <https://unfccc.int/process-and-meetings/the-paris-agreement/what-is-the-paris-agreement> Accessed on 11 November 2020.
- van Wesemael, B., Paustian, K., Andrén, O., Cerri, C.E.P., Dodd, M., Etchevers, J., Goidts, E., Grace, P., Kätterer, T., McConkey, B.G., Ogle, S., Pan, G. & Siebner, C. 2010, How can soil monitoring networks be used to improve predictions of organic carbon pool dynamics and CO<sub>2</sub> fluxes in agricultural soils? *Plant and Soil*, **338**(1-2), s. 247-259. DOI 10.1007/s11104-010-0567-z.
- Vanguelova, E.I., Bonifacio, E., Vos, D., B, Hoosbeek, M.R., Berger, T.W., Vesterdal, L., Armolaitis, K., Celi, L., Dinca, L., Kjønnaas, O.J., Pavlenda, P., Pumpanen, J., Püttsepp, Reidy, B., Simončič, P., Tobin, B. & Zhiyanski, M. 2016a, Sources of errors and uncertainties in the assessment of forest soil carbon stocks at different scales—review and recommendations, *Environmental monitoring and assessment*, **188**(11), s. 1-24. DOI 10.1007/s10661-016-5608-5.
- Vaudour, E., Bel, L., Gilliot, J.M., Coquet, Y., Hadjar, D., Cambier, P., Michelin, J. & Houot, S. 2013, Potential of SPOT Multispectral Satellite Images for Mapping Topsoil Organic Carbon Content over Peri-Urban Croplands, *Soil Science Society of America journal*, **77**(6), s. 2122-2139. DOI 10.2136/sssaj2013.02.0062.
- Verra. About us, Who We Are. <https://verra.org/about-verra/who-we-are/>. Accessed on 12 October 2020.
- Verra. VM0042 Methodology for Improved Agricultural Land Management, v1.0. 2020. <https://verra.org/methodology/vm0042-methodology-for-improved-agricultural-land-management-v1-0/>. Accessed on 27 October 2020.
- Vicharnakorn, P., Shrestha, R., Nagai, M., Salam, A. & Kiratiprayoon, S. 2014. Carbon Stock Assessment Using Remote Sensing and Forest Inventory Data in Savannakhet, Lao PDR, *Remote sensing (Basel, Switzerland)*, **6**(6), s. 5452-5479. DOI 10.3390/rs6065452.

- Vierling, K.T., Vierling, L.A., Gould, W.A., Martinuzzi, S. & Clawges, R.M. 2008, Lidar: shedding new light on habitat characterization and modeling, *Frontiers in ecology and the environment*, **6**(2), s. 90-98. DOI 10.1890/070001.
- Viscarra Rossel, R.A., Adamchuk, V.I., Sudduth, K.A., McKenzie, N.J. & Lobsey, C. 2011, *Proximal Soil Sensing: An Effective Approach for Soil Measurements in Space and Time*, Elsevier.
- Vitti, C., Stellacci, A.M., Leogrande, R., Mastrangelo, M., Cazzato, E. & Ventrella, D. 2016, Assessment of organic carbon in soils: a comparison between the Springer–Klee wet digestion and the dry combustion methods in Mediterranean soils (Southern Italy), *Catena (Giessen)*, **137**, s. 113-119. DOI 10.1016/j.catena.2015.09.001.
- Watson, C. 2009. Forest carbon accounting: overview and principles. Food and Agriculture Organization of the United Nations. [http://www.undp.org/content/dam/aplaws/publication/en/publications/environment-energy/www-ee-library/climate-change/forest-carbon-accounting-overview---principles/Forest\\_Carbon\\_Accounting\\_Overview\\_Principles.pdf](http://www.undp.org/content/dam/aplaws/publication/en/publications/environment-energy/www-ee-library/climate-change/forest-carbon-accounting-overview---principles/Forest_Carbon_Accounting_Overview_Principles.pdf). Accessed on 4 August 2020.
- Wendt, J.W. & Hauser, S. 2013, An equivalent soil mass procedure for monitoring soil organic carbon in multiple soil layers, *European Journal of Soil Science*, **64**(1), s. 58-65. DOI 10.1111/ejss.12002.
- Whitehead, D., Baisden, T., Campbell, D., Curtin, D., Davis, M.R., Hedley, C.B., Beare, M., Dr, Jones, H., Kelliher, F.M., Saggarr, S. & Schipper, L. 2012, *Review of soil carbon measurement methodologies and technologies, including nature and intensity of sampling, their uncertainties and costs*, Ministry for Primary Industries, Wellington.
- Wielopolski, L., Chatterjee, A., Mitra, S. & Lal, R. 2011, In situ determination of Soil carbon pool by inelastic neutron scattering: Comparison with dry combustion, *Geoderma*, **160**(3), s. 394-399. DOI 10.1016/j.geoderma.2010.10.009.
- Wielopolski, L., Hendrey, G., Johnsen, K.H., Mitra, S., Prior, S.A., Rogers, H.H. & Torbert, H.A. 2008, Nondestructive System for Analyzing Carbon in the Soil, *Soil Science Society of America Journal*, **72**(5), s. 1269-1277. DOI 10.2136/sssaj2007.0177.
- Wild, A. 1993. *Soils and the Environment: An Introduction*. Cambridge, Great Britain: Cambridge University Press. 287 s.
- Willgoose, G. 2018. Soils: Soil Depth. In *Principles of Soilscape and Landscape Evolution* (pp. 90-95). Cambridge: Cambridge University Press. doi:10.1017/9781139029339.007



# CARBON FARMING SCHEME

LIFE Preparatory Project

[www.st1.com/st1-life](http://www.st1.com/st1-life)

Coordinator:



Partners:



Tyynelän  
tila



puro •  
earth

NEOT  
North European Oil Trade

Financed by:



LIFE19 PRE FI001 – SI2.828588  
The Life Carbon Farming  
project has received funding  
from the LIFE Programme of  
the European Union